

Eresia phillyra (I. p. 185).

To the localities given, add:—MEXICO, Jalapa (*F. D. G.*), Orizaba (*Elwes*), Cuesta de Misantla (*Trujillo*), Atoyac (*H. H. Smith*); GUATEMALA, Sinanja in Vera Paz (*Champion*).

Eresia alsina (I. p. 186).

To the localities given, add:—NICARAGUA, Matagalpa (*Richardson*).

Eresia myia (I. p. 188).

To the localities given, add:—MEXICO, Sierra Madre de Tepic (*Richardson*), Atoyac, Teapa (*H. H. Smith*), Valladolid in Yucatan (*Gaumer*); HONDURAS, San Pedro Sula (*Wittkugel*).

Eresia ofella (I. p. 189).

To the localities given, add:—GUATEMALA, Volcan de Santa Maria (*Richardson*).

Eresia clara (I. p. 189).

To the localities given, add:—MEXICO, Teapa (*H. H. Smith*); HONDURAS, San Pedro Sula (*Wittkugel*); COSTA RICA, Santa Clara Valley (*Zurcher*); PANAMA, Bugaba (*Champion*).

ARGYNNIS (to follow the genus *Eresia*, I. p. 189).

Argynnis, Fabr. in Illiger's Mag. vi. p. 283 (1807).

1. **Argynnis nokomis**.

Argynnis nokomis, W. H. Edwards, Proc. Acad. Phil. 1862, p. 221 (♂)¹; Butt. N. Am. i., *Argynnis*, iv. tabb. (♂ ♀)²; Snyder, Occasional Memoirs of the Chicago Ent. Soc. i. p. 33 (1900)³.

♀. Alis obscure fuscis, maculis in seriebus tribus irregularibus marginibus externis subparallelis nigris, fascia communi lata, ad apicem anticarum in maculas varias fracta ad angulum analem posticarum extendente, albida, maculis ad margines externos albis, aliis nigris quatuor transversis in cellula anticarum: subtus anticis rufo-fulvis, fere dimidio exteriori pallide flavo, maculis nigris ut in pagina superiore, maculis tribus apicem versus et serie submarginali (nisi ad angulum analem) argenteis, costa viridi-ochracea; posticis quoque viridi-ochraceis, fascia lata margini externo propiore, venis divisa, pallide flava, maculis submarginalibus septem, octo in serie subparalleli interioribus, aliis basin propioribus, omnibus argenteis nigro circumcinctis.

Hab. NORTH AMERICA, Rocky Mountains¹, Bitter Root Mountains², Southern Utah³, Arizona³.—MEXICO, near Durango city (*Becker*).

Mr. Becker has sent us a single female specimen of this species from Durango. It differs from Edwards's figures of that sex of *A. nokomis*, and from an Arizonan female in our collection, having rather more than the basal half of each wing almost uniformly fuscous between the black spots, and on the secondaries beneath the interspaces between the silvery spots from the base to the middle obscure greenish ochreous