

(*F. D. G.*), Cordova (*Rümelí*), Xucumanatlan, Acaguizotla, La Venta, Venta de Zopilote (*H. H. Smith*), Acapulco (*J. J. Walker*).

We now have a very long series of this insect.

Phyciodes elada (I. p. 196).

To the localities given, add:—MEXICO, Durango city (*Becker*), Lake Chapala, Jalisco (*Richardson*), Tepetlapa, Amula, Hacienda de la Imagen and Venta de Zopilote in Guerrero (*H. H. Smith*).

We now have many examples of this species from Western Mexico.

Phyciodes imitata (I. p. 196).

To the localities given, add:—MEXICO, Northern Sonora (*Morrison*).

13 (A). **Phyciodes dymas**. (Tab. CVIII. figg. 13, 14, ♂.)

Melitæa dymas, Edw. Canad. Ent. ix. p. 190¹.

Melitæa larunda, Strecker, Lep. Rhop. & Het. p. 130².

P. imitata similis, sed anticis fascia costali ultra cellulam pallida: subtus posticis lineis nigris discalibus interruptis.

♀ mari similis.

Hab. NORTH AMERICA, California, Texas^{1 2}.—MEXICO, Northern Sonora (*Morrison*), Alamos (*Buchan-Hepburn*).

This species, as we had already anticipated, inhabits Northern Mexico, whence a long series has been sent us by Mr. Morrison from Sonora, and one from Alamos in Chihuahua by Buchan-Hepburn. It is a very small narrow-winged insect, differing from *P. imitata* as noted above. We figure one of Mr. Morrison's specimens.

13 (B). **Phyciodes hepburni**, sp. n. (Tab. CVIII. figg. 15, 16, ♂.)

P. imitata similis, sed multo obscuriore, fascia maculari discali communi (in anticis interrupta) et macula parva in cellula, pallide flavis: subtus posticis fasciis albescentioribus.

♀ mari similis.

Hab. MEXICO, Pinos Altos in Chihuahua (*Buchan-Hepburn*).

We have four specimens of this little species, including both sexes. In its narrow wings it agrees with *P. dymas* and *P. imitata*, differing from the latter as noted above. *P. elada*, which is another nearly allied form, has broader wings and the band on the underside of the secondaries less distinctly defined.

13 (C). **Phyciodes anomalus**. (Tab. CVIII. figg. 17, 18.)

Phyciodes anomalus, Godm. & Salv. Trans. Ent. Soc. 1897, p. 243¹.

Alis fusco-nigris, ciliis albo interruptis; anticis dimidio distali punctis albis notato; posticis fascia mediana fulva intra eam punctis quibusdam ejusdem coloris et extra eam punctis albis: subtus pallidioribus,