

anticis punctis albis majoribus lunulis submarginalibus saturate fulvis; posticis fascia mediana cervina extra eam lunulis nigris et punctis albis seriatim positus, lunulis fulvis submarginalibus nigro limbatis et intra eas lunulis albis, ad basin fulvo et albido maculatis, maculis omnibus late nigro marginatis; palpis fuscis, subtus albis; abdomine subtus albido supra fusco, lateribus albido punctatis.

Hab. MEXICO, Colima (*mus. Staudinger*¹).

A distinct species with no near ally. We have seen but one specimen, and that in Dr. Staudinger's collection.

16 (A). **Phyciodes faustus**. (Tab. CVIII. figg. 19, 20.)

Phyciodes faustus, Godm. & Salv. Trans. Ent. Soc. 1897, p. 243¹.

P. ptolyca similis, sed alis anticis aliter punctatis, maculis duabus medianis (inferiore majore fere rotundata), duabus subapicalibus (inferiore parva, superiore elongata), fascia mediana posticarum integra, lineola submarginali evanescente.

Hab. PANAMA, Chiriqui (*mus. Staudinger*¹).

Though compared with *P. ptolyca*, this species has only a general resemblance to that insect. The spots on the primaries are fewer in number and differently placed, only a small one remaining in the end of the cell with a larger one below it. The wings, too, are narrower.

Phyciodes fragilis (I. p. 198).

To the localities given, add:—MEXICO, San Blas (*Richardson*), Misantla, Vera Cruz (*F. D. G.*), Orizaba (*Elwes*), Atoyac, Teapa (*H. H. Smith*).

Phyciodes texana (I. p. 200).

To the localities given, add:—MEXICO, N. Sonora (*Morrison*), Pinos Altos in Chihuahua, Rio Mescales (*Buchan-Hepburn*), Ciudad Victoria, Sierra Madre de Tepic, Lake Chapala (*Richardson*), Durango city (*Becker*), Venta de Zopilote, Acaguizotla (*H. H. Smith*), Morelia, Cuernavaca, Orizaba (*F. D. G.*).

Phyciodes ptolyca (I. p. 201).

To the localities given, add:—MEXICO, Jalapa, Misantla (*F. D. G.*), Atoyac, Teapa (*H. H. Smith*); COSTA RICA, Santa Clara Valley (*Zurcher*).

24 (A). **Phyciodes phlegias**, sp. n. (Tab. CVIII. figg. 21, 22, ♂.)

P. ptolyca similis, sed anticis ad marginem externum magis erosis, maculis majoribus laete fulvis; posticis fascia transversa lata ultra cellulam lineaque angusta interrupta margini propiore coloris ejusdem: subtus posticis fascia transversa albida, introrsum costam versus litura rubro-fusca notatis, extrorsum ad marginem externum fusco-rubro suffusis.

Hab. HONDURAS (*ex Staudinger*).

We possess two males of this species. It is a close ally of *P. ptolyca*, differing as noted above.