

Phyciodes tulcis (I. p. 203).

To the localities given, add:—MEXICO, Tampico and Ciudad Victoria in Tamaulipas, Sierra Madre de Tepic, San Blas (*Richardson*), Vera Cruz (*F. D. G.*), Dos Arroyos, Atoyac, Teapa (*H. H. Smith*); GUATEMALA, Volcan de Santa Maria (*Richardson*); HONDURAS, Ruatan I. (*Gaumer*); NICARAGUA, Matagalpa (*Richardson*).

Phyciodes ardys (I. p. 204).

To the localities given, add:—MEXICO, Omilteme, Acaguizotla, Soledad, Amula (*H. H. Smith*), Orizaba (*Elwes*), Cuesta de Misantla (*Trujillo*); NICARAGUA, Matagalpa (*Richardson*).

With the long series of specimens we now possess from Mexico, we are convinced that *P. hermas* (Hew.) cannot be separated from *P. ardys*, of which *P. subota* and *P. drymæa*, G. & S., are probably only extreme forms.

32 (A). **Phyciodes alexon**. (Tab. CVIII. figg. 23, 24, ♂; 25, 26, ♀.)

Phyciodes alexon, Godm. & Salv. Ann. & Mag. Nat. Hist. (6) iii. p. 353 (♂ ♀) (1889)¹.

P. nebulosæ affinis, sed antecarum maculis omnibus pallide flavidis (nec ferrugineis), lineis et maculis dimidiis distalis posticarum quoque flavidis distinguenda.

♀ mari similis, sed major et maculis omnibus magis distinctis.

Hab. MEXICO, Cuernavaca, Rincon, and Acaguizotla in the Sierra Madre del Sur (*H. H. Smith*¹).

Described from three males and one female sent us by Mr. H. H. Smith¹. Allied to *P. nebulosa*, having elongated wings and the under surface similarly coloured.

Phyciodes argentea (I. p. 207).

To the localities given, add:—MEXICO, Atoyac (*H. H. Smith*, *Schumann*); GUATEMALA, Solola (*Richardson*).

40 (A). **Phyciodes cyno**. (Tab. CVIII. figg. 27, 28, ♂; 29, 30, ♀.)

Phyciodes cyno, Godm. & Salv. Ann. & Mag. Nat. Hist. (6) iii. p. 354 (♂) (1889)¹.

♂. *P. otani* similis, sed anticis maculis discalibus majoribus et plaga sericea nulla: subtus anticis (apice excepto) fere unicoloribus, lineolis ad basin et costam versus obsoletis vix notatis; posticis plerumque griscentioribus, maculis omnibus sicut in *P. otane*, sed magis distinctis.

♀. Feminae *P. sopoli* similis, sed maculis omnibus albis (nec fulvis), iis ad basin propioribus minutioribus; posticis disco rufo-fulvo suffusis, fascia angulata interiore multo angustiore: subtus griscentioribus, anticis maculis albis ut supra.

Hab. MEXICO, Orizaba (*H. J. Elwes*¹), Cuesta de Misantla (*coll. Schaus*).

Our original description of this species was taken from a single male specimen captured by Mr. Elwes, but we have since seen a pair of it in Mr. Schaus's collection; our figure of the female is taken from one of the latter.