

Peridromia amphinome (I. p. 271).

Ageronia amphinome, Staud. Exot. Schmett. p. 128, t. 44^e.

To the localities given, add:—MEXICO, Cuesta de Misantla (*Trujillo*), Rio Papagaio, Atoyac (*H. H. Smith*); HONDURAS, San Pedro Sula (*Wittkugel*); NICARAGUA, Matagalpa, (*Richardson*); COSTA RICA, Santa Clara Valley (*Zurcher*).

Peridromia guatemalena (I. p. 273).

To the localities given, add:—MEXICO, Tetamoa (*Buchan-Hepburn*), La Venta in Guerrero, Teapa (*H. H. Smith*); HONDURAS, San Pedro Sula (*Wittkugel*); NICARAGUA, Matagalpa (*Richardson*); COSTA RICA, Santa Clara Valley (*Zurcher*).

DIDONIS (I. p. 275).

Didonis aganisa (I. p. 276).

To the localities given, add:—MEXICO, Tampico (*Richardson*), Acaguizotla, Atoyac (*H. H. Smith*), Cuesta de Misantla, San Lorenzo (*Trujillo*); HONDURAS, San Pedro Sula (*Wittkugel*), Ruatan I. (*Gaumer*); NICARAGUA, Matagalpa (*Richardson*).

Didonis biblis (I. p. 277).

Didonis biblis, Staud. Exot. Schmett. p. 128, t. 44^e.

To the localities given, add:—PANAMA⁶, Peña Blanca (*Champion*), Panama city (*Buchan-Hepburn*).

CYSTINEURA (I. p. 277).

Cystineura amymone (I. p. 278).

To the localities given, add:—NORTH AMERICA, Texas.—MEXICO, Pinos Altos in Chihuahua (*Buchan-Hepburn*), Tampico, Sierra Madre de Tepic (*Richardson*), Guerrero, Atoyac (*H. H. Smith*), Jalapa (*F. D. G.*), Coatepec (*Brooks*); NICARAGUA, Matagalpa (*Richardson*).

We now have a specimen from Jalapa with a distinct fulvous streak at the middle of the outer margin of the primaries, as in the two Texan examples in our collection sent us by Mr. Strecker under the name of *C. dorcas*.

TIMETES (I. p. 282).

Dr. Skinner [Syn. Cat. N. Am. Rhopalocera (1898)] has recorded three of our species from the Southern United States; these are, *T. peleus*, Sulz. (= *petreus*, Cram.), from Florida, and *T. coresia*, Godt., and *T. chiron*, Fabr., from Texas.

Timetes harmonia (I. p. 286).

To the localities given, add:—MEXICO, Atoyac in Vera Cruz (*H. H. Smith*).

Sent us in numbers by Mr. Smith.