

Timetes berania (I. p. 286).

Megalura berania, Staud. Exot. Schmett. p. 135, t. 45⁴.

To the localities given, add:—HONDURAS, San Pedro Sula (*Wittkugel*); COSTA RICA, Santa Clara Valley (*Zurcher*).

Timetes merops (I. p. 288).

To the localities given, add:—COSTA RICA, Santa Clara Valley (*Zurcher*).

Timetes chiron (I. p. 288).

To the localities given, add:—MEXICO, Tampico (*Richardson*), Campala (*Forrer*), Pinal, Puebla (*Elwes*), Tierra Colorada in Guerrero (*H. H. Smith*).

Timetes coresia (I. p. 289).

Megalura coresia, Staud. Exot. Schmett. p. 134, t. 45⁴.

To the localities given, add:—MEXICO, Orizaba (*Elwes*), Cuesta de Misantla (*Trujillo*), Atoyac (*H. H. Smith*); GUATEMALA, Purula (*Champion*).

Timetes iole (I. p. 289).

To the localities given, add:—COSTA RICA, Santa Clara Valley (*Zurcher*).

PYRRHOGYRA (I. p. 291).

Pyrrhogyra hypsenor (I. p. 292).

To the localities given, add:—MEXICO, San Blas (*Richardson*); GUATEMALA, Volcan de Santa Maria (*Richardson*), Panzos (*Conradt*); HONDURAS, San Pedro Sula (*Wittkugel*).

Figured by us under the name of *P. tipha* (Tab. XXVII. figg. 3, 4).

ADELPHA (I. p. 294).

Adelpha leuceria (I. p. 300).

To the localities given, add:—MEXICO, Cuesta de Misantla (*Trujillo*); PANAMA, Chiriqui (*Trötsch*).

We have now received four specimens of this species from the Atlantic slope of Mexico.

Adelpha fessonia (I. p. 301).

To the localities given, add:—MEXICO, Sierra Madre de Tepic (*Richardson*), Jalapa (*F. D. G.*); GUATEMALA, Volcan de Santa Maria (*Richardson*); NICARAGUA, Matagalpa (*Richardson*).

Adelpha oberthüri (I. p. 302).

To the localities given, add:—HONDURAS, San Pedro Sula (*Wittkugel*).