

**Anæa electra** (I. p. 352).

*Anæa electra*, Staud. Exot. Schmett. p. 178, t. 61<sup>5</sup>.

To the localities given, add:—MEXICO, Jalapa (*F. D. G.*), Orizaba (*Elwes*); GUATEMALA, Zapote (*Champion*).

## PROTOGONIUS (I. p. 357).

**Protopogonius cecrops** (I. p. 357).

To the localities given, add:—MEXICO, Atoyac, Teapa (*H. H. Smith*); NICARAGUA, Matagalpa (*Richardson*).

## EURYGONA (I. p. 362).

1 (A). **Eurygona cheles**. (Tab. CX. figg. 1, 2, ♀.)

*Eurygona cheles*, Godm. & Salv. Ann. & Mag. Nat. Hist. (6) iii. p. 356 (♀) (1889)<sup>1</sup>.

♀. *E. aurantiacæ* similis, sed anticis area circa ramum medianum primum et cellula fusco lavatis, posticarum margine externo rotundato (haud angulato): subtus pallidioribus, linea transversa communi in anticis fere recta, in posticis magis sinuata.

*Hab.* PANAMA, Chiriqui (*Trötsch, in mus. Staudinger*<sup>1</sup>).

The single female from which our original description was taken still remains the only specimen known to us of this species.

**Eurygona argentea** (I. p. 365).

To the localities given, add:—PANAMA, Chiriqui (*mus. Staudinger*).

3 (A). **Eurygona corduena**.

*Eurygona corduena*, Hew. Boliv. Butt. p. 13<sup>1</sup>; Ex. Butt., *Erycinidæ*, ff. 4, 5, 6 (♂ ♀)<sup>2</sup>.

Alis fusco-nigris, anticis ad basin, posticis ad angulum analem læte nitide cyaneis, his ciliis ad medium marginis externi hic illuc albis: subtus argenteo-albis; anticis margine externo et fasciis tribus transversis ad costam divergentibus fuscis, una per cellulam, duabus discalibus, fasciis his in posticis continuatis, curvatis, margini externo plus minusve parallelis, in margine interno terminatis, duabus externis plaga magna rufo-aurantia interruptis, ocello nigro inter venam medianam et ramo suo secundo ornata.

♀. Alis fuscis, ad margines externos saturatioribus; anticis plaga magna ovali alba ornatis.

*Hab.* PANAMA, Volcan de Chiriqui (*Trötsch*).—PERU; BOLIVIA<sup>1 2</sup>.

We are unable to separate a male *Eurygona* submitted to us by Dr. Staudinger from *E. corduena*, described by Hewitson from specimens obtained by Buckley in Bolivia. This species is evidently allied to *E. argentea*.

**Eurygona eubule** (I. p. 366).

To the localities given, add:—MEXICO, Atoyac (*H. H. Smith*).

Mr. Smith has sent us specimens of both sexes of this species from Atoyac. The two females agree with the one figured by us.

**Eurygona sergia** (I. p. 367).

To the localities given, add:—MEXICO, Atoyac (*H. H. Smith*).