

**Mesosemia nivalis** (I. p. 386).

To the localities given, add:—MEXICO, Atoyac, Teapa (*H. H. Smith*).

## CREMNA (I. p. 387).

**Cremna umbra** (I. p. 389).

To the localities given, add:—MEXICO, Atoyac, Jalapa (*Schaus*), Teapa (*H. H. Smith*).

## DIORHINA (I. p. 390).

**Diorhina butes** (I. p. 391).

To the localities given, add:—MEXICO, Rincon in Guerrero (*H. H. Smith*), Cuesta de Misantla (*Trujillo*), Atoyac (*Schumann, H. H. Smith*), Jalapa (*coll. Schaus*).

## ERYCINA (I. p. 392).

**Erycina jurgensensi** (I. p. 393).

To the localities given, add:—MEXICO, Cordova (*coll. Schaus*).

## LEPRICORNIS (I. p. 405).

**Lepricornis melanchroia** (I. p. 405).

To the locality given, add:—MEXICO, Atoyac in Vera Cruz (*H. H. Smith*).

We have now nine examples of this species, including both sexes, from Atoyac.

1 (A). **Lepricornis unicolor**.

*Esthemopsis* (?) *unicolor*, Godm. & Salv. anteà, i. p. 487, t. 47. f. 6<sup>1</sup>.

Having a series of *L. melanchroia*, Feld., before us, we find that the insect we doubtfully referred to the genus *Esthemopsis* (anteà, i. p. 487) belongs properly to *Lepricornis*, and is indeed a close ally of Felder's species. The type of *E. unicolor* proves to be a male, not a female as we had supposed.

## LIMNAS (I. p. 408).

**Limnas pixe** (I. p. 409).

To the localities given, add:—MEXICO, Rincon, La Venta and Rio Papagaio in Guerrero, Atoyac, Teapa (*H. H. Smith*), Omealca, Cuesta de Misantla and San Lorenzo (*Trujillo*), Jalapa (*coll. Schaus*); HONDURAS, San Pedro Sula (*Wittkugel*); NICARAGUA, Matagalpa (*Richardson*).

1 (A). **Limnas gynæceas**. (Tab. CX. fig. 3, ♀.)

*Limnas gynæceas*, Godm. & Salv. Ann. & Mag. Nat. Hist. (6) iii. p. 356 (1889)<sup>1</sup>.

♀. Alis nigris; anticis apice flavidis, macula ad basin coccinea; posticis, præter maculam basalem coccineam, immaculatis: subtus ut supra, abdomine maculis quatuor coccineis utrinque notato.

*Hab.* MEXICO, Cuesta de Misantla (*M. Trujillo*<sup>1</sup>).