

We have a single female of this insect, which may prove to be only an extreme form of *L. pixe* with the scarlet marginal spots of the secondaries absent. We have since received specimens of *L. pixe* from the same locality.

**Limnas cephise** (I. p. 411).

To the localities given, add:—MEXICO, La Venta and Tierra Colorada in Guerrero (*H. H. Smith*), Jalapa (*coll. Schaus*); GUATEMALA, Volcan de Santa Maria (*Richardson*).

**Limnas acroleuca** (I. p. 412).

To the localities given, add:—MEXICO, Milpas in Durango, Ventanas (*Forrer*), Venta de Zopilote in Guerrero (*H. H. Smith*), Jalisco (*Schumann*).

SYMMACHIA (I. p. 415).

**Symmachia emesia** (I. p. 416).

To the locality given, add:—MEXICO, Mazatlan (*ex H. Edwards*).

**Symmachia yucatanensis** (I. p. 416).

To the localities given, add:—MEXICO, Venta de Zopilote in Guerrero (*H. H. Smith*), Jalapa (*coll. Schaus*).

We now question whether this is a valid species, as a specimen of *S. emesia* has been sent us from Mexico.

**Symmachia accusatrix** (I. p. 416).

To the localities given, add:—MEXICO, Atoyac (*H. H. Smith*), Jalapa (*Trujillo*).

Specimens of both sexes of this species have now been sent us from Eastern Mexico; we had not hitherto seen females from Central America.

**Symmachia championi** (I. p. 417).

To the locality given, add:—MEXICO, Jalapa (*coll. Schaus*).

**Symmachia rubina** (I. p. 419).

To the localities given, add:—MEXICO, Teapa (*H. H. Smith*).

MESENE (I. p. 420).

**Mesene croceella** (I. p. 423).

To the localities given, add:—MEXICO, Atoyac (*H. H. Smith*), Jalapa (*coll. Schaus*).

**Mesene margaretta** (I. p. 423).

To the localities given, add:—MEXICO, Atoyac (*H. H. Smith*), Cuesta de Misantla (*Trujillo*), Jalapa (*coll. Schaus*).