

CHARIS (I. p. 427).

Charis zama (I. p. 428).

To the localities given, add:—MEXICO, Atoyac (*H. H. Smith*).

Charis velutina (I. p. 428).

To the localities given, add:—MEXICO, Atoyac, Teapa (*H. H. Smith*), Omealca, San Lorenzo (*Trujillo*), Jalapa (*coll. Schaus*).

Sent us in abundance from South-east Mexico. The Teapa females differ from those from Atoyac in having a common dark discal angulated band crossing the wings.

Charis laverna (I. p. 430).

To the localities given, add:—MEXICO, Durango city (*Becker*), Chilpancingo, Rio Papagaio, Venta de Zopilote (*H. H. Smith*), Jalapa (*coll. Schaus*).

Charis holosticta (I. p. 431).

To the localities given, add:—MEXICO, Jalapa (*coll. Schaus*).

Charis sulphurea (I. p. 432). (Tab. CX. fig. 4, ♀.)

To the locality given, add:—MEXICO, Teapa (*H. H. Smith*), Jalapa (*coll. Schaus*).

CALYDNA (I. p. 433).

Calydna venusta (I. p. 433).

To the localities given, add:—MEXICO, Rincon in Guerrero (*H. H. Smith*).

Calydna hegias (I. p. 434).

To the localities given, add:—MEXICO, Tampico (*Richardson*), Chilpancingo (*H. H. Smith*), Jalapa (*coll. Schaus*).

SAROTA (I. p. 434).

Sarota chrysus (I. p. 435).

To the localities given, add:—MEXICO, Atoyac, Teapa (*H. H. Smith*), Jalapa (*coll. Schaus*).

Sarota myrtea (I. p. 437).

To the localities given, add:—MEXICO, Cordova (*Elwes*), Jalapa (*coll. Schaus*).

ANTEROS (I. p. 437).

Anteros carausius (I. p. 440).

To the localities given, add:—MEXICO, Jalisco (*Schumann*), Rincon in Guerrero, Atoyac (*H. H. Smith*), Jalapa (*coll. Schaus*); NICARAGUA, Matagalpa (*Richardson*).