

**Caria rhacotis** (I. p. 449).

To the localities given, add:—COSTA RICA, Santa Clara Valley (*Zurcher*).

## BÆOTIS (I. p. 452).

**Bæotis zonata** (I. p. 453).

To the localities given, add:—MEXICO, Rincon, Atoyac (*H. H. Smith*), Jalapa (*coll. Schaus*).

2. **Bæotis nesæa**. (Tab. CX. figg. 11, 12, ♂.)

*Bæotis nesæa*, Godm. & Salv. Ann. & Mag. Nat. Hist. (6) iii. p. 356 (♂) (1889)<sup>1</sup>.

Alis pallide flavis, anticis dimidio distali, costa et linea obliqua ad basin, nigricantibus, limbo marginali maculis duabus flavis, una subapicali, secunda ad angulum analem, macula rubra ad costæ medium; posticis margine externo late nigricante (linea flava et altera exterior plumbea includente) cum linea basali margini interno subparalleli conjuncta: subtus ut supra, sed maculis et lineis submarginalibus majoribus, lineis basalibus subobsoletis.

*Hab.* PANAMA, Chiriqui (*Trötsch, in mus. Staudinger*<sup>1</sup>).

Dr. Staudinger has submitted to us the single male specimen here figured and described. It has no near allies known to us, but somewhat resembles *B. felix*, Hew. The red spot, however, on the costa of the primaries is quite characteristic.

## METACHARIS (I. p. 453).

2. **Metacharis (?) melucina**.

*Metacharis (?) melucina*, Staud. Exot. Schmett. p. 256, t. 91<sup>1</sup> (*erythromelas* ♂, nec ♀).

Alis nigris, anticis plaga magna infra venam medianam, ultra cellulam et ad marginem internum extendente, cum maculis duabus in cellula confluentibus, aurantiis; posticis margine costali albo, litura a basi (extrorsum latiore) haud ad marginem externum extendente aurantia, anticis et posticis linea angusta nitente plumbea submarginali: subtus anticis fuscis, area infra venam medianam pallide ochracea, costa et maculis duabus (una costali ultra cellulam, secunda majore exterior) maculisque quibusdam angulum analem versus, albis; posticis albis, nigro maculatis.

*Hab.* PANAMA, Chiriqui<sup>1</sup> (*Ribbe, ex Staudinger*). — COLOMBIA, San Pablo, Rio San Juan<sup>1</sup>.

Dr. Staudinger has sent us a male of this species from Chiriqui, and two others of the same sex from the Rio San Juan, under the above name, an example from the latter locality having been figured by him as *M. erythromelas* (Sepp). It is a close ally of that insect, which inhabits the Amazon Valley, but differs in having the orange patch on both wings more extended. The Chiriqui specimen, from which the above description is taken, differs again from the southern insects in having two orange spots in the cell confluent with the discal patch. These species have been placed by Dr. Staudinger doubtfully in *Metacharis*, from which they will probably have to be removed. The two insects have very much the facies of *Lemonias emylius* (Cram.).