

12 (A). **Theope eleutho.** (Tab. CX. figg. 29, 30, ♀.)

Theope eleutho, Godm. & Salv. Trans. Ent. Soc. 1897, p. 247¹.

♀. Alis fusco-nigris; anticis ad basin viridescenti-cæruleis, costa omnino nigricante, maculis tribus subapicalibus in linea obliqua viridescenti-cæruleis; posticis area interna et area discali quoque viridescenti-cæruleis, hac venis nigris intersecta, costa late nigricante, margine interno albicante: subtus glauco-fuscescentibus, linea communi ab angulo anticarum apicali ad medium marginis interni posticarum saturate fusca, anticis ad basin rufescentibus, maculis duabus in cellula albis, posticis maculis parvis septem submarginalibus nigris, alia alba ad cellulæ basin; corpore supra cæruleo, subtus alis concoloribus.

Hab. PANAMA, Chiriqui (*Trötsch, in mus. Staudinger*¹).

Dr. Staudinger has sent us a female of this species, which differs from all others of the genus in our collection. It belongs to the section having a common line on the underside running from the apical angle of the primaries to the middle of the inner border of the secondaries. The apical angle of the primaries is acute and salient, owing to the outer margin being concave towards the point; the anal angle, too, of the secondaries is acute.

12 (B). **Theope mania.** (Tab. CX. figg. 31, 32, ♀.)

Theope mania, Godm. & Salv. Trans. Ent. Soc. 1897, p. 247¹.

♀. *T. eleuthonti* similis, sed alis cærulescentioribus, anticarum apice minus acuto; posticis (præter costam nigram) omnino cæruleis, angulo anali minus producto: subtus pallidioribus, anticis ad basin vix rufescentibus, posticis maculis submarginalibus evanescentibus.

Hab. MEXICO, San Blas (*Richardson*¹).

This is a species having a dark line across the under surface of both wings. We have only received a single female, taken by Mr. Richardson in Western Mexico.

12 (C). **Theope bacenis.**

Theope bacenis, Schaus, Ent. Amer. vi. p. 19 (♂) (1890)¹.

Alis anticis nigris, margine interno ad basin late cæruleo; posticis cæruleis, margine costali nigro: subtus brunneis, fascia communi ab angulo anticarum apicali ad medium marginis analis posticarum saturate fusca, anticis ad basin flavescentibus.

Hab. MEXICO, Coatepec¹ (*coll. Schaus*).

15. **Theope matuta.** (Tab. CX. figg. 33, 34, ♀.)

Theope matuta, Godm. & Salv. Trans. Ent. Soc. 1897, p. 246¹.

♀. Alis nigris, anticis ad basin (costa nigra excepta) nitide cæruleis, venis divisis; posticis area interna usque ad basin nitide cærulea, costa late, margine externo angustiore, nigris: subtus fuscis, anticis costa ad basin rufa, maculis elongatis discalibus albidis transfasciatis, fascia maculosa submarginali coloris ejusdem; posticis maculis sagittiformibus; fronte inter oculos nigra, utrinque albido marginata; palpis nigris antice albidis; corpore fusco, abdomine linea mediana alba.

Hab. COSTA RICA, Santa Clara Valley 1200 feet (*Zurcher*¹).

Allied to *T. thebais* of the Amazon Valley, but with the blue on the upperside of