

the wings much brighter. Beneath, the wings are darker and the pale markings more definite, and the rufous base of the costa of the primaries is more extended. We have seen only a single female of this distinct species.

ISAPIS (I. p. 485).

**Isapis hera** (I. p. 486).

To the locality given, add:—GUATEMALA, Panzos (*Conradt*).

EUMÆUS (II. p. 5).

**Eumæus minyas** (II. p. 6).

To the localities given, add:—MEXICO, La Venta, Rincon, Atoyac, Teapa (*H. H. Smith*); GUATEMALA, Volcan de Santa Maria (*Richardson*); HONDURAS, San Pedro Sula (*Wittkugel*); COSTA RICA, Santa Clara Valley (*Zurcher*).

**Eumæus debora** (II. p. 7).

To the localities given, add:—MEXICO, Ciudad Victoria in Tamaulipas (*Richardson*), Atoyac (*H. H. Smith*), Orizaba (*Elwes*).

THECLA (II. p. 8).

**Thecla regalis** (II. p. 12).

To the localities given, add:—MEXICO, Cuesta de Misantla (*Trujillo*), Atoyac (*H. H. Smith*); HONDURAS, Ruatan I. (*Gaumer*); NICARAGUA, Matagalpa (*Richardson*).

**Thecla cypria** (II. p. 13).

To the localities given, add:—MEXICO, Rincon in Guerrero (*H. H. Smith*).

**Thecla damo** (II. p. 15).

To the localities given, add:—MEXICO, Tierra Colorada, Acapulco, Rincon, La Venta, Teapa (*H. H. Smith*), Atoyac (*Schumann*), Cuesta de Misantla (*Trujillo*); HONDURAS, San Pedro Sula (*Wittkugel*); NICARAGUA, Matagalpa (*Richardson*); COSTA RICA, Santa Clara Valley (*Zurcher*).

**Thecla augustula** (II. p. 16).

To the localities given, add:—MEXICO, Cuesta de Misantla (*Trujillo*).

We now have a pair of this insect from Mexico.

**Thecla nepia** (II. p. 16).

To the localities given, add:—MEXICO, Acaguizotla, Atoyac, Teapa (*H. H. Smith*); NICARAGUA, Matagalpa (*Richardson*); PANAMA, Chiriqui (*Trötsch, ex Staudinger*).