

Thecla ortygnus (II. p. 41).

To the localities given, add:—GUATEMALA, Volcan de Santa Maria (*Richardson*).

Thecla jebus (II. p. 42).

To the localities given, add:—MEXICO, Acaguizotla in Guerrero (*H. H. Smith*).

Thecla brescia (II. p. 43).

To the localities given, add:—MEXICO, Bolaños in Jalisco (*Richardson*), Tierra Colorada, Rincon, Venta de Zopilote, Dos Arroyos, Atoyac (*H. H. Smith*); NICARAGUA, Matagalpa (*Richardson*).

Thecla ligurina (II. p. 44).

To the localities given, add:—GUATEMALA, Volcan de Santa Maria (*Richardson*).

Thecla lyde (II. p. 44).

To the localities given, add:—MEXICO, Atoyac, Teapa (*H. H. Smith*).

Thecla temesa (II. p. 45).

To the localities given, add:—MEXICO, Teapa (*H. H. Smith*).

This insect now proves to extend as far north as South-east Mexico, whence Mr. Smith has sent us a single male example.

Thecla mycon (II. p. 46).

To the localities given, add:—MEXICO, Rincon, Atoyac (*H. H. Smith*).

Thecla phobe (II. p. 47).

To the localities given, add:—MEXICO, Atoyac (*H. H. Smith*).

Thecla xami (II. p. 48).

♀. Alis fulvis, marginibus externis et costa anticarum nigro-fuscis: subtus sicut in mari.

To the localities given, add:—MEXICO, Amula (*H. H. Smith*).

84 (A). **Thecla spinetorum**. (Tab. CXI. figg. 5, 6, ♂.)

Thecla spinetorum, Hew. Ill. Diurn. Lep. p. 94, t. 45. ff. 198, 199¹; Boisd. Ann. Soc. Ent. Belg. xii. p. 42².

Thecla ninus, Edw. Trans. Am. Ent. Soc. iii. p. 270³.

Alis et corpore supra fuscis, schistaceo-cæruleo lavatis: subtus rufo-ferrugineis, linea communi transversa, introrsum nigro limbata et in posticis marginem internum versus valde angulata, alba; posticis serie duplici macularum nigrarum ad marginem externum, exteriore maculis duabus ad angulum analem proximis magis distinctis et albo notatis.

Hab. NORTH AMERICA, Colorado³, California^{1 2}.—MEXICO, Bolaños in Jalisco (*Richardson*).