

This species must be added to the Central-American list, as we possess a worn specimen of it from Mexico, which agrees with an example from California in our collection. We are indebted to Dr. Holland for its identification, and he has been kind enough to compare a drawing of our Californian insect, which we figure, with Edwards's type of *T. ninus* from Colorado.

Thecla ocrisia (II. p. 49).

To the localities given, add :—MEXICO, Tampico, Lake Chapala (*Richardson*), Atoyac (*H. H. Smith*); NICARAGUA, Matagalpa (*Richardson*).

Thecla keila (II. p. 50).

To the localities given, add :—MEXICO, Acaguizotla in Guerrero (*H. H. Smith*).

Thecla tephraeus (II. p. 51).

To the localities given, add :—MEXICO, Acaguizotla, Venta de Zopilote, Dos Arroyos, Rincon, Atoyac, Teapa (*H. H. Smith*).

Thecla volupia (II. p. 53).

♀. Alis murinis vix caeruleo basin versus suffusis: subtus sicut in mari.

To the localities given, add :—GUATEMALA, Polochic Valley (*F. D. G., O. S.*).

Since the publication of the preceding pages we have detected the female of this species in our collection.

Thecla syncellus (II. p. 54).

To the localities given, add :—MEXICO, Acapulco, Rio Papagaio, Rincon, Dos Arroyos, Tierra Colorada, Atoyac, Teapa (*H. H. Smith*); GUATEMALA, Volcan de Santa Maria (*Richardson*); HONDURAS, San Pedro Sula (*Wittkugel*); NICARAGUA, Matagalpa (*Richardson*); COSTA RICA, Santa Clara Valley (*Zurcher*).

Thecla ochus (II. p. 55).

To the localities given, add :—MEXICO, Cuesta de Misantla (*Trujillo*); PANAMA, Chiriqui (*Trötsch, ex Staudinger*).

Thecla pion (II. p. 56).

To the localities given, add :—MEXICO, Orizaba (*Elwes*).

Thecla orasus (II. p. 59).

To the localities given, add :—MEXICO, Jalapa (*F. D. G.*).

Thecla quaderna (II. p. 60).

To the localities given, add :—MEXICO, Puebla 7000 feet (*Elwes*).