

3. *Philampelus typhon*.

Sphinx typhon, Klug, Neue Schmett. Mus. Berl. Heft i. t. 3. f. 1¹.

Philampelus typhon, Walk. Cat. viii. p. 177².

Hab. MEXICO^{1 2}, Oaxaca (*Deppe, mus. Oxf.*).

Of this fine species I have only seen the specimen in the Hope collection at Oxford; its nearest ally is *P. linnei*.

4. *Philampelus pandorus*.

Daphnis pandorus, Hübn. Samml. ex. Schmett. ii. t. 161. f. 3, 4.

Philampelus pandorus, Walk. Cat. viii. p. 174¹.

Hab. NORTH AMERICA¹.—MEXICO, Cordova (*Rümel*); GUATEMALA, Coban (*Champion*); PANAMA, Chiriqui (*Ribbé*).—ANTILLES, Jamaica (*mus. D.*).

A very variable species, closely allied to *P. lycaon* (Cr.).

5. *Philampelus satellitia*.

Sphinx satellitia, Linn. Mant. i. p. 539¹; Drury, Ill. i. t. 29. f. 1².

Philampelus satellitia, Butl. Trans. Z. S. p. 576³.

Hab. BRITISH HONDURAS³; NICARAGUA, Chontales (*Belt*); PANAMA, Chiriqui (*Ribbé, Arcé, mus. D.*).—ANTILLES, Jamaica³; COLOMBIA, Antioqua (*mus. D.*); ECUADOR (*mus. D.*).

This species is the largest of the genus; its nearest ally is *P. lycaon*.

6. *Philampelus anchemolus*.

Sphinx anchemolus, Cr. Pap. Ex. t. 224. f. C¹.

Philampelus anchemolus, Walk. Cat. viii. p. 178²; Butl. Trans. Z. S. ix. p. 577³.

Hab. MEXICO, Mazatlan (*H. Edwards*); PANAMA, Chiriqui (*mus. Staudinger*).—GUIANA, Surinam^{1 2}; RIO JANEIRO³.

Very closely allied to *P. satellitia*.

7. *Philampelus labruscæ*.

Sphinx labruscæ, Linn. Mus. Lud. Ulr. p. 352; Clerck, Icones Ins. t. 47. f. 3.

Philampelus labruscæ, Walk. Cat. viii. p. 178¹.

Hab. MEXICO¹ (*Boucard, mus. D.*).—ANTILLES, Jamaica¹, San Domingo¹; CUBA (*mus. D.*); COLOMBIA¹; ECUADOR (*mus. D.*); VENEZUELA¹; ARGENTINE REPUBLIC (*mus. D.*).

This species occurs in Mexico; but as yet I have not seen a specimen from any other