

4. *Ambulyx sexoculata*.

Ambulyx sexoculata, Grote, Ann. Lyc. N. Y. viii. p. 204¹.

Hab. GUATEMALA (*mus. Brit.*).—BRAZIL¹.

This species is allied to *A. rostralis* and *A. gannascus*.

Subfam. *SMERINTHINÆ*.

SMERINTHUS.

Smerinthus, Latreille, Hist. Nat. Ins. iii. p. 431 (1802).

About six species are placed in this genus, two being found in Central America.

1. *Smerinthus ophthalmicus*.

Smerinthus ophthalmicus, Boisd. Ann. Soc. Ent. Belge, xii. p. 67¹; Strecker, Lep. Rhop. & Het. pt. 7, t. 7. ff. 4, 5²; Butl. Trans. Z. S. ix. p. 592³.

Hab. N. AMERICA, California².—MEXICO^{1 3} (*mus. Brit.*).

2. *Smerinthus saliceti*. (Tab. I. fig. 9.)

Smerinthus saliceti, Boisd. Sp. Gen. des Lép. Hét. p. 35¹.

Hab. MEXICO¹.

I have only seen the type specimen of this species, kindly lent me by Mons. C. Oberthür to figure. It is nearly allied to *S. ophthalmicus*, but a much smaller species.

Subfam. *SPHINGINÆ*.

AMPHONYX.

Amphonyx, Poey, Cent. Lep. Cuba (1832).

This genus is closely allied to *Acherontia*, from which it mostly differs in having longer and narrower wings, more slender antennæ, and semitransparent hind wings.

1. *Amphonyx duponcheli*.

Amphonyx duponcheli, Poey, Cent. Lep. Cuba¹; Butl. Trans. Z. S. ix. p. 599².

Macrosilia antæus (part.), Walk. Cat. viii. p. 200³.

Hab. PANAMA, Chiriqui (*Arcé, mus. D.*).—ANTILLES, Jamaica^{2 3}, San Domingo^{2 3}, Cuba¹ (*mus. D.*), Trinidad (*mus. D.*).

I have not seen specimens of this species from any locality north of Panama.

2. *Amphonyx medor*.

Sphinx medor, Cr. Pap. Ex. iv. t. 394. f. A¹.

Amphonyx medor, Butl. Trans. Z. S. ix. p. 599².

Hab. MEXICO², Jalapa (*Höge*), Valladolid, Yucatan (*Gaumer*).—ECUADOR (*mus. D.*); GUIANA, Surinam¹.

Allied to the preceding species, but altogether darker in colour.