

2. *Ægeria senta*, sp. n. (Tab. V. fig. 1.)

Primaries hyaline, the costal margin, discal mark, and outer margin black; secondaries hyaline, the fringe black: head, thorax, and abdomen black; antennæ black; palpi white beneath, black above; collar white; legs black, with tufts of blackish hair on the tibiæ. Expanse $\frac{1}{2}$ inch.

Hab. GUATEMALA, Panzos (*Champion*).

A very small species, not nearly allied to any with which I am acquainted.

3. *Ægeria sellustiformis*, sp. n. (Tab. V. fig. 2.)

Primaries whitish hyaline, the costal margin and the apex black, the discal mark black; secondaries with the fringe only black: head, antennæ, and thorax blue-black; palpi white beneath, black above; legs black; tarsi banded with white. Expanse $\frac{3}{4}$ inch.

Hab. PANAMA, Bugaba 800 to 1500 feet (*Champion*).

This species is allied to *Æ. producta*, but is a much larger insect.

4. *Ægeria ruficaudis*. (Tab. V. fig. 6.)

Ægeria ruficaudis, Walk. Cat. xxxi. p. 6¹.

Sesia ruficaudis, Boisd. Sp. Gén. des Lép.-Hét. i. p. 451².

Hab. PANAMA, San Feliz (*Champion*).—UPPER AMAZONS, Ega (*Bates*^{1 2}).

The single specimen sent by Mr. Champion agrees well with Walker's type, but not with his description, as far as the wings are concerned.

5. *Ægeria deipyla*, sp. n. (Tab. V. fig. 8.)

Primaries hyaline, with a golden-yellow shade, the costal margin and the apex brown; discal band brown, edged with yellow, inner margin yellow; secondaries hyaline, the outer margin slightly banded with brown: head and thorax black, banded with yellow; palpi lemon-yellow; antennæ black above, beneath (excepting at the tips) reddish brown; abdomen black, with all the segments banded with yellow; the anal tuft black, tipped with white; beneath, the abdomen is bright yellow; legs pale yellow. Expanse $\frac{3}{4}$ inch.

Hab. PANAMA, Bugaba, 800 to 1500 feet (*Champion*).

This species is not unlike *Æ. ichneumoniformis*, but differs in many respects.

6. *Ægeria producta*. (Tab. V. fig. 7.)

Ægeria producta, Walk. Cat. xxxi. p. 4¹.

Sesia producta, Boisd. Sp. Gén. des Lép.-Hét. i. p. 449.

Hab. MEXICO¹, Oaxaca (*Sallé*); PANAMA, Bugaba 800 to 1500 feet (*Champion*).

The type specimen is in poor condition. The species is allied to *Æ. rhyssæformis*, but still very distinct. The figure is taken from a Bugaba specimen.

7. *Ægeria læta*. (Tab. V. fig. 10.)

Ægeria læta, Walk. Cat. viii. p. 59.

Sesia læta, Boisd. Sp. Gén. des Lép.-Hét. i. p. 459.

Hab. GUATEMALA, San Gerónimo (*Champion*).

Walker's type is a very faded specimen, and without any locality; but it is without doubt the same as the Guatemala examples before me, one of which I have figured.