

8. *Ægeria aurata*.*Ægeria aurata*, H. Edwards, Papilio, i. p. 190.*Hab.* PANAMA (*H. Edwards*).Allied to *Æ. ceriiformis*, Led.9. *Ægeria flava*.*Ægeria flava*, H. Edwards, Papilio, i. p. 189.*Hab.* PANAMA, Colon (*Mead*).Size of *Æ. tipuliformis*.10. *Ægeria guatemalena*, sp. n. (Tab. V. fig. 9.)

Hyaline: primaries slightly shaded with yellow near the base; the costal margin and apex brown; secondaries hyaline with the fringe brown in some lights; the wings are quite opalescent: head, palpi, and antennæ pale yellow beneath, above black; abdomen black, banded with yellow, the whole of the under surface pale yellow; legs yellow. Expanse  $1\frac{1}{10}$  inch.

*Hab.* GUATEMALA, San Gerónimo (*Champion*).

This species is allied to *Æ. cynipiformis* (Esp.), but is easily distinguished from that species by the opalescent colour of the wings, and the greater number of yellow bands on the abdomen.

11. *Ægeria tabogana*, sp. n. (Tab. V. fig. 16.)

Primaries dark brown, yellowish hyaline on the inner margin from the base to near the anal angle; secondaries hyaline, the outer margin slightly brown: head, thorax, and abdomen black, the back of thorax and the abdomen banded with yellow; antennæ black; palpi pale yellow; collar yellow; legs black above, yellowish white on the underside. Expanse  $\frac{3}{4}$  inch.

*Hab.* PANAMA, Taboga Island (*Champion*).

Mr. Champion only obtained a single specimen of this very distinct species.

12. *Ægeria ceres*, sp. n. (Tab. V. fig. 11.)

Primaries glossy black, greenish at the base, a small round hyaline spot beyond the cell near the anal angle; secondaries hyaline, the outer margin and fringe black: head, palpi, thorax, and abdomen black; anal tuft black; legs black, tarsi banded with yellow. Expanse  $\frac{3}{4}$  inch.

*Hab.* GUATEMALA, Totonicapan 8500 to 10,500 feet (*Champion*).

This species is very distinct. Only a single specimen has been sent.

13. *Ægeria proserpina*, sp. n. (Tab. V. fig. 13.)

Primaries glossy greenish black; secondaries dull black, the basal third hyaline: head, palpi, and antennæ black, the collar and underside of thorax reddish brown; upperside of thorax and all the legs black; abdomen wanting. Expanse  $\frac{3}{4}$  inch.

*Hab.* GUATEMALA, San Gerónimo (*Champion*).

A small species, very distinct from any with which I am acquainted. Only a single specimen was received.