

2. *Eupyra corydon*, sp. n. (Tab. VI. fig. 13.)

Primaries glossy black, thickly speckled with green scales; a narrow green band crossing the wing close to the base; two small hyaline spots about the middle of the cell, three at the end, second and third very minute, and four beyond nearer the apex; secondaries black, with a central band of four bluish hyaline spots, the fourth below the others and close to the apex: head, thorax, and abdomen black, two white spots on each side of the thorax; antennæ black, rather deeply pectinated; legs black. Expanse $1\frac{3}{4}$ inch.

Hab. GUATEMALA, Senahu (*Champion*).

This appears to be a very scarce insect. Mr. Champion only obtained a single specimen. Dr. Staudinger has sent me a specimen also from Guatemala, but without the exact locality.

CALONOTOS.

Calonotos, Hübner, Verz. bek. Schmett. p. 123; Walk. Cat. i. p. 234.

Mr. Butler has divided this genus into three sections; but I am not able to see his reason for so doing; the insects placed in these sections differ considerably, and I should have thought them worthy of generic distinction; but not having a sufficient number of specimens I am unable to determine this point.

1. *Calonotos nexa*.

Læmocharis nexa, Herr.-Schäff. Samml. aussereur. Schmett. f. 254; Walk. Cat. vii. p. 1627.

Calonotos nexa, Butl. Journ. Linn. Soc., Zool. xii. p. 368¹.

Hab. MEXICO, Valladolid, Yucatan (*Gaumer*); GUATEMALA, Cubilguitz (*Champion*).—UPPER AMAZONS¹.

The specimens of this insect from Mexico are the smallest I have seen; but in other respects they do not differ in any way.

2. *Calonotos tiburtus*.

Sphinx tiburtus, Cram. Pap. Exot. iii. t. 237. f. C¹.

Calonotos tiburtus, Hübn. Verz. bek. Schmett. p. 12.

Euchromia tiburtus, Walk. Cat. i. p. 244.

Hab. PANAMA, Bugaba 800 to 1500 feet (*Champion*); Volcan de Chiriqui (*Trötsch, mus Staudinger*).—GUIANA, Surinam¹.

I have specimens of both sexes before me; they agree well with Cramer's figure, being closely allied to *C. nexa*.

3. *Calonotos interrupta*.

Calonota interrupta, Walk. Cat. vii. p. 1628¹.

Hab. MEXICO¹.

4. *Calonotos trotschi*, sp. n. (Tab. VIII. fig. 8.)

Primaries black, crossed beyond the middle by a band of four hyaline spots, the first and second close to the costal margin quite small, the third and fourth larger, three elongated streaks near the base bluish hyaline; secondaries black, with the base and a small spot near the apex hyaline: head black, with a