

Glaucopis (Isanthrene) flavicornis, Walk. Cat. i. p. 155 (nec Fabr.).

Isanthrene incendiaria, Butl. Journ. Linn. Soc., Zool. xii. p. 375.

Hab. MEXICO, Milpas (*Forrer*).—BRAZIL (*mus. D.*).

A single example of this species was sent; it agrees well with specimens in my own collection from South Brazil.

5. *Isanthrene crabroniformis*. (Tab. VI. fig. 23.)

Isanthrene crabroniformis, Staudinger, Verh. z.-b. Ges. Wien, xxv. p. 120 (1875)¹.

Hab. PANAMA, Volcan de Chiriqui and David (*Ribbe*¹, *Champion*), Colon (*Boucard*, *mus. D.*), Taboga Island (*Champion*).

A very common species in Chiriqui, most nearly allied to *I. flavicornis*, Fabr., from Colombia.

HOMŒOCERA.

Homœocera, Felder, Reise der Nov. Lep. t. 102. f. 26.

The type of this genus is *Homœocera crassa*, Felder. Mr. Butler in his paper on the Zygænidæ includes eight species in this genus, three of which occur in our country.

1. *Homœocera jansonis*.

Gymnelia jansonis, Butl. & Druce, Cist. Ent. i. p. 116.

Gymnelia jansonis, Butl. Lep. Exot. t. 61. f. 17¹.

Homœocera jansonis, Butl. Journ. Linn. Soc., Zool. xii. p. 375.

Hab. COSTA RICA, Cartago¹ (*Van Patten*), Irazu (*Rogers*); PANAMA, Bugaba (*Champion*, *Ribbe*, *mus. Staudinger*).

I have only seen three examples of this species; the type is now in the British Museum.

2. *Homœocera beatrix*, sp. n. (Tab. VI. fig. 25.)

Primaries and secondaries yellowish hyaline, the outer and inner margins bordered with black, the base of the primaries black, spotted with bright green scales: head and thorax black, banded and spotted with bright green; abdomen black, broadly banded with bright metallic green, each segment narrowly margined with pale yellow; anus bright orange; antennæ black, very slightly pectinated; legs brownish black. Expanse 1½ inch.

Hab. PANAMA, Volcan de Chiriqui (*Ribbe*), Bugaba and San Feliz (*Champion*).

This beautiful little species is allied to *H. jansonis*, from which it differs in having extremely narrow margins to the primaries, and being of a much more yellow colour.

3. *Homœocera scintillans*.

Læmocharis scintillans, Herr.-Schäff. Samml. aussereur. Schm. f. 244.

Glaucopis (Pæcilosoma) flavitarsis, Walk. Cat. i. p. 158¹.

Homœocera scintillans, Butl. Journ. Linn. Soc., Zool. xii. p. 376; Ill. Lep. Het. i. p. 26, t. 12. f. 4.

Hab. COSTA RICA (*Van Patten*).—VENEZUELA¹.

A single example in bad condition was received in Van Patten's last collection; it is