

2. *Gymnelia xenodorus*, sp. n. (Tab. VII. fig. 1.)

Primaries and secondaries yellowish hyaline, the outer and inner margins broadly banded with black, a black band at the end of the cell: head and collar blue; antennæ black, tipped with white; thorax black; abdomen black, shot with blue on the sides, the first segment dirty white, the fifth, sixth, and seventh orange-red on the upperside; legs black. Expanse $1\frac{1}{2}$ inch.

Hab. GUATEMALA, Panzos (*Champion*).

This species is allied to *G. whitelyi*, from Eastern Peru; but it is at once distinguished from that insect by its bright-red extremity of the abdomen.

LÆMOCHARIS.

Læmocharis, Herrich-Schäffer (part), Samml. aussereur. Schmett. 258.

This genus, as restricted by Mr. Butler, now has about eight species placed in it, all of them being small. Herrich-Schäffer included a large number of species in his genus, many of them belonging to quite different genera. Three species occur in our country, two of them being new.

1. *Læmocharis trigutta*.

Glaucopis (Pseudomya) trigutta, Walk. Cat. i. p. 145¹.

Læmocharis trigutta, Butl. Journ. Linn. Soc., Zool. xii. p. 383; Ill. Lep. Het. i. p. 29, t. 7. f. 10.

Hab. GUATEMALA, Panzos (*Champion*).—COLOMBIA¹; AMAZONS, Tabatinga (*M. Stuart, mus. D.*).

This species is found over a very extended range of country. The example from Guatemala is darker in colour than those from South America, but in other respects it does not differ.

2. *Læmocharis stryma*, sp. n. (Tab. VII. fig. 5.)

Primaries and secondaries hyaline, with the veins and outer margins black: head and antennæ black; thorax black, with a small spot on each side and one at the base bright carmine; abdomen black; legs black, the tarsus of the hind legs white. Expanse $\frac{3}{4}$ inch.

Hab. NICARAGUA, Chontales (*Belt*).

3. *Læmocharis hercyna*, sp. n. (Tab. VII. fig. 6.)

Primaries hyaline, the base, inner and outer margins narrowly bordered with black, the apex broadly black; secondaries hyaline, with the outer margin black: head, thorax, and antennæ black; abdomen pale chrome-yellow; legs black. Expanse 1 inch.

Hab. NICARAGUA, Chontales (*Belt*); PANAMA, Bugaba (*Champion*).

A pretty little species, resembling *Ilipa stilbosticta*, Butler, from Colombia.

GYMNOPODA.

Gymnopoda, Felder, Reise d. Nov. Lep. t. 102. f. 22; Index, p. 5.

Dr. Felder states that this genus is allied to *Horama*. The primaries are long and narrow, as in that genus, the palpi are rather long, and the antennæ slightly pectinated.