

1. *Gymnopoda subflamma*, sp. n. (Tab. VIII. fig. 10.)

Primaries semitransparent black, except from the base to the end of the cell, which is quite hyaline, almost the same as *Gymnopoda ochracea*, Felder; secondaries hyaline, the outer margin and apex dusky black: the head and antennæ black, collar with a few whitish scales; thorax and legs black; abdomen reddish yellow, with a central band of black spots; the anus black. Expanse 1 inch.

Hab. PANAMA, Chiriqui (*Champion*).

Of this insect I have only received a single example. It is closely allied to Dr. Felder's species, from which it chiefly differs by having a white collar and a row of black spots down the centre of the abdomen.

THRINACIA.

Thrinacia, Butler, Journ. Linn. Soc., Zool. xii. p. 384 (1876).

Two species are placed in this genus by Mr. Butler, one of which occurs in our country, also two new species now described.

1. *Thrinacia afflicta*.

Glaucopis (Pseudomya) afflicta, Walk. Cat. i. p. 144¹.

Thrinacia afflicta, Butl. Journ. Linn. Soc., Zool. xii. p. 384; Ill. Lep. Het. i. p. 29, t. 7. f. 12.

Hab. PANAMA, Chiriqui (*Champion*).—BRAZIL¹.

2. *Thrinacia sanguinea*, sp. n. (Tab. VII. fig. 9.)

Primaries black, hyaline in the centre of the wing; secondaries black: head, antennæ, and abdomen black; thorax and the base of the primaries bright red; legs black. Expanse 1 inch.

Hab. GUATEMALA, Teleman (*Champion*).

This species differs from *T. afflicta*, Walk., in having the primaries more hyaline and the base of the wings all red.

3. *Thrinacia intermedia*, sp. n.

Closely allied to the preceding species, the chief difference being its more sooty colour, red instead of black head, slightly more pectinated antennæ, longer and narrower primaries, the abdomen and legs black. Expanse 1 inch.

Hab. PANAMA, Bugaba 8000 feet (*Champion*).

A very distinct species, of which a single example was sent.

PSEUDOCHARIS, gen. nov.

General appearance of *Læmocharis*; neurulation of the primaries as in *Dycladia*; secondaries very small and narrow at the apex. Head small; palpi minute, first joint oval, second elongated, thickly clothed with scales. Antennæ very slightly pectinated, with thirty-nine joints. Thorax and abdomen short and thick, the same as in *Dycladia*. Fore leg long and slender, the coxa very flat; femur long and slight; tibia rather short, with a short spur rather beyond the middle; tarsus five-jointed, hind leg very long, tibia with four spurs, the first two about the middle, the second pair at the end close to the tarsus, which