

The three female specimens before me are very similar in colour and markings; the one from Mexico in Mr. Schaus's collection is paler than the others and has the dusky lines across the wings less distinct; the one from Guatemala is considerably darker. The male is apparently unknown.

## 2. *Drepanodes moneta*, sp. n. (Tab. XLIV. fig. 2, ♀.)

*Female*. Primaries and secondaries pale brownish-fawn-colour; the primaries crossed from the apex to the inner margin by a narrow dark brown line, edged on the inner side with a very narrow silvery-white line, a curved dark fawn-coloured line near the base, a second line crossing the wing from the costal margin beyond the cell and joining the line extending from the apex to the inner margin beyond the middle, the apex and the outer margin thickly irrorated with black scales; the secondaries crossed about the middle by a continuation of the line on the primaries, and with a waved dark fawn-coloured submarginal line extending from the apex to the anal angle, and a black dot in the middle of the cell; the fringe reddish-brown; the underside pale yellowish-fawn-colour, thickly irrorated with small black spots, the outer margins of both wings broadly bordered with dark reddish-brown: head, antennæ, thorax, abdomen, and legs pale fawn-colour. Expanse  $1\frac{1}{2}$  inch.

*Hab.* MEXICO, Coatepec (*coll. Schaus*).

Apparently allied to *D. pionaria*, Walk., but very distinct from it.

## 3. *Drepanodes falcularia*.

*Patalene falcularia*, Herr.-Schäff. Samml. aussereur. Schmett. p. 80, fig. 199<sup>1</sup>.

*Comibana falcularia*, Herr.-Schäff. loc. cit.<sup>2</sup>

*Drepanodes falcularia*, Guen. Sp. gén. des Lép. ix. p. 69<sup>3</sup>; Walk. Cat. xx. p. 77<sup>4</sup>.

*Hab.* MEXICO, Paso de San Juan in Vera Cruz (*coll. Schaus*); GUATEMALA, Volcan de Atitlan 2500 to 3500 feet (*Champion*); PANAMA, Bugaba 800 to 1500 feet (*Champion*).—GUIANA, Surinam<sup>2 3 4</sup>.

This species varies very much on the upperside: in some specimens the line crossing both wings is dark brown, in others it is almost wanting; the yellowish-white spots near the anal angle of the primaries, as shown in Herrich-Schäffer's figure, are sometimes entirely obliterated or replaced by blackish patches or dots. The underside shows little variation.

## 4. *Drepanodes astanda*, sp. n. (Tab. XLIV. fig. 3.)

Primaries and secondaries orange-yellow; the primaries slightly clouded at the base with a darker colour, and with a black dot in the middle of the cell, and a faint reddish-brown line extending from the apex to the inner margin above the anal angle, the outer margin of the wing irrorated with silvery-white scales; the secondaries with a small black dot in the cell and two indistinct blackish-brown patches below the middle nearest the anal angle, the outer margin irrorated with silvery-white scales; the underside of both wings bright yellow, the outer margins broadly bordered with dark reddish-brown: head, thorax, antennæ, and legs pale yellow; (abdomen wanting). Expanse  $1\frac{1}{4}$  inch.

*Hab.* MEXICO, Teapa in Tabasco (*H. H. Smith*).

One specimen. Allied to *D. falcularia*, Herr.-Schäff.