

edge of the costal margin close to the apex white,—that of the secondaries orange-yellow, irrorated along the costal margin with brown, and with a black dot at the end of the cell: head, thorax, and abdomen above yellowish-olive-colour, beneath and the legs orange-yellow, the antennæ brown. Expanse $1\frac{3}{4}$ inch.

Hab. PANAMA, Chiriqui (*Trötsch, in mus. Staudinger*).

One specimen.

6. **Acrotomia minucia**, sp. n. (Tab. XLV. figg. 21, ♂; 22, ♀.)

Male. Primaries and secondaries pale brownish-grey, irrorated with grey; the primaries crossed from the apex to the middle of the inner margin by a reddish-brown line, this line extending across the middle of the secondaries, both wings with a submarginal waved white line, the fringe grey; the underside pale yellow, irrorated with brown scales, both wings with a rather wide submarginal dark brown line (which is not nearly so broad in some specimens as in others), the primaries with three small white spots close to the apex: head, thorax, abdomen, and legs pale fawn-colour, the antennæ brown.—*Female.* Primaries and secondaries pinkish-fawn-colour, slightly irrorated with brown scales, both wings crossed from the costal to the inner margin by a wide greyish band, which is thickly irrorated with black, and has black lines on either side, the band on the primaries extending from the middle of the wing to the apex and widened towards the costal and inner margins, the fringe fawn-colour; the underside very similar to that of the male, but in some specimens the submarginal brown band on the secondaries is wanting. Expanse, ♂ ♀, $1\frac{5}{16}$ inch.

Hab. MEXICO, Cordova (*Rümelì*), Coatepec (*coll. Schaus, Brooks*), Atoyac in Vera Cruz (*H. H. Smith*); PANAMA, Chiriqui (*Ribbe, in mus. Staudinger*).—ECUADOR.

This insect varies considerably on the underside, not only in colour but in the extent of the submarginal brown band.

We figure a pair from Chiriqui.

7. **Acrotomia (?) bola**, sp. n. (Tab. XLV. fig. 23, ♂.)

Male. Primaries and secondaries of a uniform mouse-colour, the secondaries slightly darker round the outer margin, both wings crossed by two fine waved brown lines, the fringe brown; the underside reddish-brown, the primaries crossed from the apex to the inner margin by a dark brown line: head, thorax, abdomen, antennæ, and legs mouse-colour. Expanse $1\frac{5}{16}$ inch.

Hab. PANAMA, Chiriqui (*Trötsch, in mus. Staudinger*).

One specimen.

8. **Acrotomia (?) nisa**, sp. n. (Tab. XLV. fig. 24, ♂.)

Male. Primaries and secondaries very pale fawn-colour, crossed from the costal to the inner margin by a wide pale yellow band, which is edged on either side with a fine brown line, and with a submarginal waved brown line, the secondaries with a very distinct black spot at the end of the cell, the fringe yellowish-brown; the underside pale brownish-white, the primaries with a submarginal waved brown line: head, thorax, abdomen, and legs pale fawn-colour, the antennæ pale brown. Expanse $1\frac{1}{2}$ inch.

Hab. GUATEMALA, San Gerónimó (*Champion*).

Two specimens.