

Hab. MEXICO, Amecameca (*F. D. G.*).

One specimen, captured by Mr. Godman in April 1888.

4. *Therina seminudata*.

Ellopiia seminudata, Walk. Cat. xxvi. p. 1508 (σ)¹.

Therina seminudaria, Packard, in Report of the U. S. Geol. Survey of the Territ. x. p. 495, t. 12. figg. 4, 5 (1876)².

Ellopiia siccaria, Walk. Cat. xxxv. p. 1547³.

Ellopiia bibularia, Grote & Robins. Ann. Lyc. N. York, viii. p. 455, t. 15 A. fig. 8⁴.

Ellopiia pellucidaria, Grote & Robins. loc. cit. p. 456, t. 15 A. fig. 9⁵.

Hab. NORTH AMERICA³, Canada^{1 2}, United States^{2 4 5}.—MEXICO, Jalapa, Coatepec (*coll. Schaus*); GUATEMALA, Las Mercedes 3000 feet (*Champion*).

We are now able to trace this common North-American species southwards into Mexico and Guatemala.

A male from Jalapa agrees well with Dr. Packard's fig. 4 (σ). Apparently a rare insect within our limits, as we have only received two specimens.

5. *Therina bada*, sp. n. (Tab. XLVI. figg. 1, σ ; 2, φ .)

Male. Primaries dull brown, the secondaries paler in colour; the primaries crossed from the costal to the inner margin by two pale yellowish waved lines, and with a black dot at the end of the cell, the fringe brown; the secondaries crossed about the middle by a very faint pale line; the underside considerably paler in colour than above: head, thorax, abdomen, antennæ, and legs dull brown. The female very similar to the male, but with the primaries mottled with pale yellowish-brown. Expanse, σ $1\frac{1}{2}$, φ $1\frac{3}{4}$ inch.

Hab. GUATEMALA, Totonicapam 8500 to 10,500 feet, Quiche Mountains 7000 to 9000 feet (*Champion*).

Varies in colour from dull brown to reddish-fawn-colour. In some male specimens the lines crossing the wings are much more distinct than in others; the females are generally paler than the males, and are still more variable in colour.

6. *Therina* (?) *nomia*, sp. n. (Tab. XLVI. fig. 4, σ .)

Male. Primaries pale purplish-brown, crossed about the middle from the costal to the inner margin by two waved dark brown lines, which are faintly edged with yellow, and with a waved submarginal line extending from the apex to the anal angle, and a small brown spot at the end of the cell; the secondaries paler in colour, and crossed about the middle from the costal to the inner margin by a dark brown waved line; the fringe of both wings purplish-brown; the underside much paler than above, and entirely without markings: head, antennæ, and thorax purplish-brown, the underside of the latter, the abdomen, and legs yellowish-brown. Expanse $1\frac{3}{4}$ inch.

Hab. MEXICO, Jalapa (*Höge*), Las Vigas (*coll. Schaus*).

In Mr. Schaus's collection there are three males and a female of this species. We have only received one specimen. A male from Las Vigas is figured.