

RICA, Volcan de Irazu 6000 to 7000 feet (*Rogers*); PANAMA, Volcan de Chiriqui 2000 to 3000 feet (*Champion*).—CUBA, San Cristobal.

We have received a good series of this species from Guatemala. *L. meganira* is allied to *L. ablinearia*, from which it may be easily distinguished by the entirely different markings of the wings; the greyish lines, however, are much more distinct in some specimens than they are in others. Our description and figure are taken from a Cahabon specimen.

### CABERODES.

*Caberodes*, Guenée, Sp. gén. des Lép. ix. p. 135 (1857); Walker, Cat. xx. p. 166.

#### 1. *Caberodes myandaria*.

*Ellopiia myandaria*, Walk. Cat. xx. p. 164<sup>1</sup>.

*Caberodes myandaria*, Packard, in Report of the U. S. Geol. Survey of the Territ. x. p. 497<sup>2</sup>.

*Hab.* MEXICO<sup>2</sup>, Oaxaca (*Sallé*<sup>1</sup>), Jalapa (*coll. Schaus*).

One specimen, a male, in Mr. Schaus's collection, is all I have seen of this species.

#### 2. *Caberodes* — ?

*Hab.* MEXICO, Pinos Altos in Chihuahua (*Buchan-Hepburn*).

One worn specimen, apparently allied to *C. confusaria* (Hübner).

#### 3. *Caberodes axona*, sp. n. (Tab. XLVI. fig. 6, ♂.)

*Male*. Primaries and secondaries uniform reddish-fawn-colour, both wings crossed about the middle from the costal to the inner margin by a narrow dark brown line, which is edged on the outer side with yellow, the primaries with a small black dot at the end of the cell; the underside considerably paler than above, without markings of any kind: head, thorax, abdomen, legs, and antennæ reddish-fawn-colour. Expanse  $1\frac{3}{4}$  inch.

*Hab.* MEXICO, Las Vigas (*coll. Schaus*).

One specimen. Allied to *C. confusaria*, from which it differs by its much darker colour, and by the primaries not having a line across their under surface.

#### 4. *Caberodes messene*, sp. n. (Tab. XLVI. fig. 7, ♂.)

*Male*. Primaries and secondaries pale greyish-yellow, both wings crossed by a brown line, the primaries also with an angular line crossing near the base and a brown spot at the end of the cell, the fringe pale greyish-yellow; the underside paler than above, the primaries slightly irrorated with brown scales, and with the costal margin tinged with yellow: head, thorax, abdomen, and legs pale greyish-yellow; the antennæ pale brown. Expanse  $1\frac{1}{4}$  inch.

*Hab.* MEXICO, Jalapa (*M. Trujillo*).

One specimen.