

Four examples, all females. They vary considerably in colour, and the Mexican specimen corresponds best with the one from Panama. The specimen figured is from Totonicapam; it is darker in colour than any of the others.

EROSINA.

Erosina, Guenée, Sp. gén. des Lép. ix. p. 149 (1857); Walker, Cat. xx. p. 180.

A single species from Brazil was included in *Erosina* by Guenée and Walker; I now add two others to it, one of which ranges from Mexico to Peru.

1. *Erosina rusticata*.

Sabulodes rusticata (Maassen), Reiss & Stübel, Reisen in Süd-Amerika, pp. 57, 153, t. 8. fig. 1¹.

Sabulodes proximata, Dognin, Le Nat. 1891, p. 223².

Hab. MEXICO, Coatepec (*coll. Schaus*); COSTA RICA, Volcan de Irazu 6000 to 7000 feet (*Rogers*); PANAMA, Chiriqui (*Arcé, in mus. D.*), Bugaba 800 to 1000 feet, Volcan de Chiriqui 3000 to 4000 feet (*Champion*).—ECUADOR¹, Loja²; PERU.

This species is closely allied to *E. hyberniata*, Guen., the type of the genus *Erosina*. *E. hyberniata*, however, can always be distinguished from *E. rusticata* by its smaller size and generally darker colour; it is well figured by Guenée. I am indebted to M. Dognin for specimens of his *E. proximata*.

2. *Erosina meropia*, sp. n. (Tab. XLVI. fig. 10, ♂.)

Male. Primaries brownish-fawn-colour, crossed from the costal to the inner margin near the base by a faint brown line, and with a dark brown line extending from near the apex to the middle of the inner margin, the wing on the inner side of this line appearing in some lights to be shaded with pinkish-white, the outer margin also shaded with the same colour; secondaries very pale fawn-colour, considerably darker round the outer margin; the fringe of both wings pale fawn-colour; the underside of the primaries pale silky-fawn-colour, darker at the apex and along the costal margin, that of the secondaries darker and slightly irrorated with dark brown scales: head, antennæ, and thorax brownish-fawn-colour, the abdomen and legs pale fawn-colour. The female with the secondaries considerably darker than in the male. Expanse, ♂ $2\frac{1}{10}$, ♀ $2\frac{1}{4}$ inches.

Hab. COSTA RICA, Volcan de Irazu 6000 to 7000 feet (*Rogers*); PANAMA, Chiriqui (*Trötsch, in mus. Staudinger*), Volcan de Chiriqui 3000 to 4000 feet (*Champion*).

Allied to an undescribed species from Colombia in my own collection. A male from Chiriqui is figured.

SELENIA.

Selenia, Hübner, Verz. bek. Schmett. p. 292 (1816); Guenée, Sp. gén. des Lép. ix. p. 150; Walker, Cat. xx. p. 181.

Walker included six species in this genus—three from Europe, two from North America, and one from New Zealand, showing that it has a very wide distribution. Eight are now added from Mexico or Guatemala.