

5. **Selenia nabdalsa**, sp. n. (Tab. XLVI. fig. 21, ♂.)

Male. Primaries and secondaries pale brown, irrorated with slightly darker scales, a small black spot at the end of the cell on the primaries; the underside very much as above, but paler in colour: head, thorax, abdomen, antennæ, and legs pale brown. Expanse $1\frac{1}{2}$ inch.

Hab. MEXICO, Las Vigas (*coll. Schaus*).

One specimen.

6. **Selenia narcæa**, sp. n. (Tab. XLVI. figg. 17, 18, ♂.)

Male. Primaries reddish-fawn-colour, the basal two-thirds much darker, with a narrow waved line crossing from near the apex to about the middle of the inner margin, a waved line crossing near the base, a dark reddish-brown spot on the costal margin near the apex, and a submarginal indistinct waved whitish line extending from the apex to the anal angle, the outer portion of the wing shaded with brown, the fringe brown; the secondaries pale fawn-colour, with a narrow whitish line partly crossing from the inner margin above the anal angle, the wing above the line thickly irrorated with reddish-brown scales, the fringe brown; the underside reddish-fawn-colour, thickly irrorated with reddish-brown scales, the outer margin edged with grey: head, thorax, and abdomen pale brown, the thorax and abdomen beneath and the legs reddish-brown, the antennæ pale brown. Expanse $1\frac{1}{2}$ inch.

Hab. MEXICO, Las Vigas (*coll. Schaus*).

Four examples, all males.

7. **Selenia belisama**, sp. n. (Tab. XLVI. figg. 19, 20, ♂.)

Male. Primaries pale brown, slightly irrorated with grey scales, a silvery-grey curved line extending from the costal margin near the apex to the inner margin, the wing on the outer side of this line much shaded with silvery-grey (in some lights with a pink tinge), the fringe brown; secondaries very pale greyish-white, thickly irrorated with brown scales, a fine reddish-brown line crossing the wing below the middle, the outer margin and part of the inner one shaded with brown; the underside pale brown, thickly irrorated with darker brown scales: head, thorax, abdomen, and legs pale brown, the antennæ dark brown. Expanse $1\frac{3}{4}$ inch.

Hab. MEXICO, Las Vigas (*coll. Schaus*).

Four specimens, all males. Allied to *S. narcæa*.

8. **Selenia azamora**, sp. n. (Tab. XLVI. figg. 22, 23, ♂.)

Male. Primaries pale brown, with three short darker brown streaks along the costal margin, a rather wide pale fawn-coloured band crossing from the apex to the inner margin, and a narrow waved line crossing near the base, the outer margin shaded with brown; secondaries very pale silky-fawn-colour, thickly irrorated with dark brown scales, and partly crossed from the inner margin above the anal angle by two faint waved lines; the fringe of both wings pale brown; the underside pale silky-fawn-colour, thickly irrorated with orange-yellow scales: head, thorax, abdomen, antennæ, and legs pale brown. Expanse $1\frac{3}{4}$ inch.

Hab. MEXICO, Las Vigas (*coll. Schaus*).

Two specimens, both males.