

(*Gabb, in mus. D.*); PANAMA, Bugaba 800 to 1000 feet, Volcan de Chiriqui 2000 to 3000 feet (*Champion*).

Varies in colour from almost white to pale yellow. *H. nemora* is allied to an undescribed species from Ecuador in my own collection. We figure a male from Bugaba and a female from the Volcan de Chiriqui.

AZELINA.

Azelina, Guenée, Sp. gén. des Lép. ix. p. 156 (1857); Walker, Cat. xx. p. 185; Butler, Ann. & Mag. Nat. Hist. 5th ser. viii. p. 29.

Pero, Herrich-Schäffer, Samml. aussereur. Schmett. p. 63.

Mr. Butler in his paper above quoted includes fifty-four species in this genus, all but three (or four) of which are from America, where the genus ranges from New York to Brazil; and others have since been described by M. Oberthür. In our region *Azelina* is represented by twenty-three species.

1. *Azelina stuposaria*.

Azelina stuposaria, Guen. Sp. gén. des Lép. ix. p. 160 (♀)¹; Walk. Cat. xx. p. 192²; Butl. Ann. & Mag. Nat. Hist. 5th ser. viii. p. 31³.

Azelina fuscularia, Feld. & Rogenh. Reise der Novara, Lep. t. 123. fig. 11 (♀)⁴.

Hab. MEXICO, Jalapa (*coll. Schaus*); GUATEMALA (*Conradt, in mus. Staudinger*), Cerro Zunil 4000 to 5000 feet, Volcan de Atitlan 2500 to 3500 feet (*Champion*); PANAMA, Volcan de Chiriqui 2000 to 3000 feet (*Champion*).—BRAZIL^{1 2 4}, Rio Janeiro³.

Felder and Rogenhofer's figure is much redder than any specimen I have seen, but the markings are identical; it is possible, however, that *A. fuscularia* may prove to be specifically distinct.

A female from Jalapa in Mr. Schaus's collection is of a dark black-brown colour on the upperside, coming closest to those from the State of Panama; on the underside it agrees with them.

2. *Azelina saturata*.

Azelina saturata, Walk. Journ. Linn. Soc., Zool. ix. p. 196 (♀)¹; Butl. Ann. & Mag. Nat. Hist. 5th ser. viii. p. 32² (nec Feld. & Rogenh.).

Azelina emmaria, Oberth. Études d'Ent. vii. p. 22, t. 2. fig. 3³.

Hab. GUATEMALA, in the city (*Rodriguez*); COSTA RICA, Volcan de Irazu 6000 to 7000 feet (*Rogers*); PANAMA, Chiriqui (*Trötsch, in mus. Staudinger*).—COLOMBIA, Bogota^{1 2}, Antioquia; ECUADOR; PERU, Huambo³.

Walker's type of *A. saturata* is in my own collection.

The Venezuelan insect figured under this name by Felder and Rogenhofer belongs to a different species, and I propose to call it *A. rogenhoferi*.