

### 3. *Azelina honestaria*.

*Azelina honestaria*, Walk. Cat. xx. p. 258 (♂)<sup>1</sup>; Butl. Ann. & Mag. Nat. Hist. 5th ser. viii. p. 33<sup>2</sup>.

*Hab.* NORTH AMERICA, Massachusetts<sup>2</sup>.—COSTA RICA, Volcan de Irazu 6000 to 7000 feet (*Rogers*).

Our Costa Rican specimens are almost identical with Walker's type in the National Museum; the locality from whence this specimen was obtained is unknown.

### 4. *Azelina xylonaria*.

*Azelina xylonaria*, Oberth. Études d'Ent. vii. p. 24, t. 2. fig. 7 (1883)<sup>1</sup>.

*Hab.* PANAMA, Chiriqui (*Arcé, in mus. D.*).—PERU, Huambo<sup>1</sup>.

One specimen from Chiriqui, agreeing well with M. Oberthür's figure.

### 5. *Azelina hübneraria*.

*Gonodontis ancetaria*, Hübn. Samml. exot. Schmett. ii. t. 217. figg. 1, 2 (♂), 3, 4 (♀)<sup>1</sup> (nec Cram.).

*Azelina hübneraria*, Guen. Sp. gén. des Lép. ix. p. 159<sup>2</sup>; Walk. Cat. xx. p. 186<sup>3</sup>; Butl. Ann. & Mag. Nat. Hist. 5th ser. viii. p. 33<sup>4</sup>; Pack. Proc. Bost. Soc. Nat. Hist. xiii. p. 386<sup>5</sup>.

*Hab.* NORTH AMERICA, New York<sup>3 4</sup>, California<sup>5</sup>.—GUATEMALA, Cerro Zunil 4000 feet, Volcan de Atitlan 2500 to 3500 feet (*Champion*); COSTA RICA (*Gabb, in mus. D.*); PANAMA, Chiriqui (*Trötsch, in mus. Staudinger*), Volcan de Chiriqui 2500 to 3500 feet (*Champion*), Panama city (*J. J. Walker*).—ECUADOR.

Varies considerably in colour and markings. Central-American specimens are usually darker than those from the United States.

### 6. *Azelina rectisectaria*.

*Pero rectisectaria*, Herr.-Schäff. Samml. aussereur. Schmett. p. 63, fig. 325<sup>1</sup>.

*Gonodontis rectisectaria*, Herr.-Schäff. loc. cit. p. 81<sup>2</sup>.

*Azelina rectisectaria*, Guen. Sp. gén. des Lép. ix. p. 157<sup>3</sup>; Walk. Cat. xx. p. 190<sup>4</sup>; Butl. Ann. & Mag. Nat. Hist. 5th ser. viii. p. 33<sup>5</sup>.

*Hab.* MEXICO, Paso de San Juan in Vera Cruz (*coll. Schaus*); COSTA RICA (*Gabb, in mus. D.*).—ECUADOR; BRAZIL<sup>1 3 4 5</sup>.

All the examples I have seen of this species are a little brighter in colour than Herrich-Schäffer's figure. In some specimens the lines on the underside of the wings are very distinct, in others they are scarcely visible.

### 7. *Azelina hædularia*.

*Azelina hædularia*, Guen. Sp. gén. des Lép. ix. p. 158 (♂ ♀)<sup>1</sup>; Walk. Cat. xx. p. 190<sup>2</sup>; Butl. Ann. & Mag. Nat. Hist. 5th ser. viii. p. 35<sup>3</sup>.

*Hab.* MEXICO, Coatepec (*coll. Schaus*); GUATEMALA, Volcan de Atitlan 2500 to 3500