

de San Juan (*coll. Schaus*), Teapa in Tabasco (*H. H. Smith*); PANAMA, Chiriqui (*Trötsch, in mus. Staudinger*).

Mr. Schaus informs me that this is the commonest species of the genus in Mexico. The single specimen received from the Tres Marias Islands is in very poor condition, but I believe it belongs to *A. astapa*. We figure a pair from Presidio de Mazatlan.

12. ***Azelina melissa***, sp. n. (Tab. XLVII. fig. 12, ♂.)

Male. Primaries with their basal two-thirds reddish-fawn-colour, the outer portion considerably paler, two dark brown slightly waved lines crossing from the costal to the inner margin, and a small hyaline streak at the end of the cell; secondaries reddish-fawn-colour, crossed beyond the middle from the apex to the anal angle by a very fine whitish line, and with several small black dots on the outer margin close to the anal angle; the underside greyish-fawn-colour, irrorated with darker scales, the lines very faint: head, thorax, abdomen, antennæ, and legs reddish-fawn-colour. Expanse $1\frac{1}{2}$ inch.

Hab. MEXICO, Jalapa (*coll. Schaus*).

One specimen.

13. ***Azelina*** —?

Hab. BRITISH HONDURAS, Belize (*Blancaneaux*).

One specimen, apparently belonging to a distinct species, but not in sufficiently good condition for determination.

14. ***Azelina metella***, sp. n. (Tab. XLVII. fig. 13.)

Primaries and secondaries pale brown, the primaries crossed by two dark brown lines—the first near the base, the second beyond the middle, the latter edged on the inner side with reddish-brown,—the secondaries crossed about the middle by a very fine darker brown line; the underside pale brown, irrorated with dark brown scales, the lines very faint, a rather large black spot at the end of the cell on the secondaries: head, thorax, abdomen, antennæ, and legs pale brown. Expanse $1\frac{2}{10}$ inch.

Hab. PANAMA, Volcan de Chiriqui 3000 feet (*Champion*).

One specimen.

15. ***Azelina aurunca***, sp. n. (Tab. XLVII. fig. 14.)

Male. Primaries and secondaries dark brown; the primaries crossed from the costal to the inner margin by two darker brown waved lines—the first near the base, the second beyond the middle,—and with a small hyaline spot at the end of the cell, beyond which is a reddish-brown mark, the fringe dark brown; the secondaries tinged with reddish-brown at the anal angle; the underside very much paler than above, with a large black spot at the end of the cell on the secondaries. Expanse $1\frac{1}{10}$ inch.

Hab. PANAMA, Chiriqui (*Arcé, in mus. D.*; *Ribbe, in mus. Staudinger*).

This small species is allied to *A. minima*, Butl., from South-east Brazil. Two specimens, both males.

16. ***Azelina clysiaria***.

Azelina clysiaria, Feld. & Rogenh. Reise der Novara, Lep. t. 123. fig. 12 (♀)¹; Butl. Ann. & Mag. Nat. Hist. 5th ser. viii. p. 43², and Trans. Ent. Soc. Lond. 1881, p. 325³.