

23. *Azelina odonaria*.

Azelina odonaria, Oberth. Études d'Ent. vii. p. 26, t. 1. fig. 5¹.

Hab. COSTA RICA, Volcan de Irazu 6000 to 7000 feet (*Rogers*); PANAMA, Chiriqui (*Trötsch, in mus. Staudinger*).—COLOMBIA¹, Antioquia.

Specimens of this species are also contained in the National Museum, where they are erroneously labelled *Meticulodes spongiaria*, Guen.

METICULODES.

Meticulodes, Guenée, Sp. gén. des Lép. ix. p. 192 (1857); Walker, Cat. xxi. p. 295.

The two described species of this genus are from Brazil; and *Azelina xylinaria*, Guen., as noted by Mr. Butler, belongs here.

1. *Meticulodes xylinaria*.

Azelina xylinaria, Guen. Sp. gén. des Lép. ix. p. 162 (♂)¹; Walk. Cat. xx. p. 193².

Meticulodes xylinaria, Butl. Ann. & Mag. Nat. Hist. 5th ser. viii. p. 30³.

Hab. MEXICO, Jalapa (*coll. Schaus*); GUATEMALA, San Gerónimo (*Champion*); COSTA RICA, Volcan de Irazu 6000 to 7000 feet (*Rogers*); PANAMA, Bugaba (*Champion*).—BRAZIL^{1 2}.

Varies considerably in colour, some specimens being much paler than others; the females are generally darker than the males.

ODONTOPERA.

Odontopera, Stephens, Ill. Brit. Ent., Haust. iii. p. 162 (1829); Guenée, Sp. gén. des Lép. ix. p. 164; Walker, Cat. xx. p. 197.

Odontopera, if the species referred to it are really congeneric, possesses a most remarkable distribution—Europe, Abyssinia, the Cape of Good Hope, and Brazil each having one or more representatives.

Gonodontis (?) *semilutearia*, Feld. & Rogenh., the female only of which is known, seems to me best placed here, as it agrees with *O. bidentata* (Linn.), the type of the genus, in form and neuration, though very different in colour. One new species from Central America is now added.

1. *Odontopera semilutearia*.

Gonodontis (?) *semilutearia*, Feld. & Rogenh. Reise der Novara, Lep. t. 122. fig. 6 (♀)¹.

Hab. GUATEMALA, Volcan de Atitlan 2500 feet (*Champion*).—COLOMBIA, Bogota¹.

One female specimen from Guatemala, agreeing well with the figure of Felder and Rogenhofer.