

5. **Bronchelia detexta.** (Tab. XLIX. fig. 3, ♂.)

Bronchelia detexta, Walk. Cat. xxi. p. 452 (♂)¹.

Hab. MEXICO, Cordova (*Rümeli*), Jalapa, Paso de San Juan (*coll. Schaus*); PANAMA, Chiriqui (*Trötsch, in mus. Staudinger*).—ANTILLES, San Domingo¹, Jamaica¹.

Central American specimens agree very nearly with Walker's types in the National Collection. A male from Cordova is figured.

6. **Bronchelia puellaria.**

Bronchelia puellaria, Guen. Sp. gén. des Lép. ix. p. 287 (♂ ♀)¹; Walk. Cat. xxi. p. 455².

Bronchelia mundaria, Walk. Cat. xxi. p. 456 (♂)³.

Hab. MEXICO, Cuesta de Misantla (*M. Trujillo*); PANAMA, Chiriqui (*Arcé, in mus. D.; Trötsch, in mus. Staudinger*).—COLOMBIA^{1 2}; GUIANA, Cayenne^{1 2}; AMAZONS³; BRAZIL³.

I have seen four specimens of this species from Central America, three males and one female. It may readily be distinguished from all the others of the genus by the clear white ground-colour of the wings. Walker² erroneously quotes *B. conjugaria*, Guen., as a synonym.

7. **Bronchelia diffundaria.**

Bronchelia diffundaria, Walk. Cat. xxi. p. 457 (♂ ♀)¹.

Bronchelia anonaria, Feld. & Rogenh. Reise der Novara, Lep. t. 125. fig. 20².

Hab. COSTA RICA, Rio Sucio (*Rogers*); PANAMA, Volcan de Chiriqui 3000 to 4000 feet (*Champion*).—COLOMBIA²; VENEZUELA^{1 2}.

Costa Rican and Panama specimens agree perfectly with others from Honda and Merida in my own collection, and also with the type in the British Museum.

BONATEA, gen. nov.

Male. Head large; thorax rather broad; abdomen long and stout, but not extending beyond the wings. Palpi thick, porrect, the third joint somewhat elongate, very blunt at the tip. Antennæ long, filiform. Legs long, rather stout; hind tibiæ much thickened, with a very long pencil of hairs arising from the base on the inner side, the spurs small. Wings ample: primaries slightly falcate at the tip, the costal margin almost straight from the base to the apex, where it is a little rounded, the inner margin straight, the outer margin convex, feebly dentate; secondaries large, their costal margin curved, the inner margin straight, clothed with long hairs, the outer margin very convex, rather deeply dentate. Type *Oxydia duciata*, Maassen.

I include *Bonatea* in the Boarminæ with very much doubt, but I do not know where else to place it.

1. **Bonatea duciata.** (Tab. XLIX. figg. 6, 7, ♂.)

Oxydia duciata, Maassen, in Reiss und Stübel's Reisen in Süd-Amerika, p. 154, t. 7. fig. 22¹.

Hab. PANAMA, Chiriqui (*Trötsch, in mus. Staudinger*).—ECUADOR¹.