

ORTHODES (I. p. 288).

4. *Orthodes albipuncta*.

Orthodes albipuncta, Schaus, Trans. Am. Ent. Soc. xxi. p. 237 (1894)¹.

Hab. MEXICO, Jalapa (*coll. Schaus*¹).

5. *Orthodes iole*.

Orthodes iole, Schaus, Trans. Am. Ent. Soc. xxi. p. 237 (1894)¹.

Hab. MEXICO, Jalapa (*coll. Schaus*¹).

AMATHES (to follow the genus *Orthodes*, I. p. 289).

Amathes, Hübner, Verz. bek. Schmett. p. 222 (1816).

1. *Amathes temperata*. (Tab. XCIV. fig. 10.)

Amathes temperata, Schaus, Trans. Am. Ent. Soc. xxi. p. 226 (1894)¹.

Hab. MEXICO, Jalapa (*coll. Schaus*¹; *M. Trujillo*); COSTA RICA, Candelaria Mts. (*Underwood*).

We figure a Costa Rican specimen.

XANTHIA (to follow the genus *Mesogona*, I. p. 290).

Xanthia, Guenée, Ann. Soc. Ent. Fr. viii. p. 490 (1839); Sp. gén. des Léop. v. p. 389; Walker, Cat. x. p. 457.

1. *Xanthia alala*. (Tab. XCIV. fig. 11.)

Xanthia alala, Druce, P. Z. S. 1890, p. 514¹.

Male. Primaries orange-brown, shaded with pale yellow, with a round spot in the cell and a large oval spot at the end of it pale yellowish-brown, a submarginal row of small spots extending from the apex to the anal angle, the outer margin and fringe brownish; secondaries brownish-white, darker round the outer margin: head and thorax yellowish-brown, the abdomen much paler; the antennæ and legs pale brown.—*Female*. Similar to the male, but darker. Expanse $1\frac{1}{2}$ inch.

Hab. MEXICO, Mexico city (*Höge*¹); COSTA RICA, La Uruca, San José (*coll. Dognin*).

2. *Xanthia alcandra*. (Tab. XCIV. fig. 12.)

Xanthia alcandra, Druce, P. Z. S. 1890, p. 515¹.

Male. Primaries pale citron-yellow, shaded with dark brown at the base and along the costal margin to beyond the middle, with two waved lines crossing the wing from the costal to the inner margin—the first near the base, the second beyond the end of the cell,—and a large round pale brown spot about the middle of the cell and an oval similarly coloured spot at the end of it; secondaries creamy-white; the underside of both wings pale yellow: head, thorax, antennæ, and legs brownish-yellow, the abdomen yellowish-white. Expanse $1\frac{1}{2}$ inch.

Hab. MEXICO, Tierra Colorada in Guerrero 2000 feet (*H. H. Smith*¹).

One specimen.