

ANTHÆCIA (I. p. 300).

4. *Anthæcia jaguarina*.

Anthæcia jaguarina, Guen. Sp. gén. des Lép. vi. p. 184, Noctuélites, t. 9. fig. 11¹; Walk. Cat. xi. p. 694².

Hab. NORTH AMERICA^{1 2}.—MEXICO, Guadalajara in Jalisco (*Goldsmith, in mus. D.*).

We have received one specimen of this species from Mexico. Guenée's figure is a very poor one.

CIRRHOPHANUS (to follow the genus *Anthæcia*, I. p. 301).

Cirrhophanus, Grote, Canad. Ent. iv. p. 187 (1872); xv. p. 76 (1883).

1. *Cirrhophanus triangulifer*. (Tab. XCIV. fig. 16.)

Cirrhophanus triangulifer, Grote, Canad. Ent. iv. p. 187¹; xiv. p. 183²; xv. p. 76 (1883)³; Proc. Acad. Phil. 1875, p. 421, fig.⁴; J. B. Smith, Cat. Noct. Bor.-Am. (Bull. U.S. Nat. Mus. no. 44) p. 261 (1893)⁵.

Chariclea triangulifera, Grote, Illustr. Essay Noct. N. Am. p. 60, t. 3. fig. 26⁶; New Check-List N. Am. Moths, p. 36 (1882)⁷.

Chariclea pretiosa, Morrison, Proc. Bost. Soc. Nat. Hist. xviii. p. 122 (1875)⁸.

Hab. NORTH AMERICA^{1 2 7}, Kansas^{4 8}, Middle and Central States to Kansas⁵, Ohio^{3 6}, Missouri^{3 6}.—MEXICO, Durango city (*Becker*).

We have received one specimen of this species from Durango.

GROTELLA (to precede the genus *Adisura*, I. p. 301).

Grotella, Harvey, Bull. Buff. Soc. Nat. Sci. ii. p. 278 (1875).

1. *Grotella salacon*. (Tab. XCIV. fig. 18.)

Grotella salacon, Druce, Ann. & Mag. Nat. Hist. (6) xvi. p. 39 (1895)¹.

Male. Primaries glossy chrome-yellow; secondaries pale primrose-yellow, with the apex, outer margin, and fringe chrome-yellow; the underside of both wings pale glossy yellow: head, thorax, and tegulæ deep chrome-yellow; the abdomen pale yellow, the anus dark chrome-yellow; the antennæ black.—*Female*. Very similar to the male. Expanse 1 inch.

Hab. MEXICO, Tierra Colorada in Guerrero 2000 feet (*H. H. Smith*¹).

2. *Grotella pyronæa*. (Tab. XCIV. fig. 19.)

Grotella pyronæa, Druce, Ann. & Mag. Nat. Hist. (6) xvi. p. 38 (1895)¹.

Primaries chalk-white, crossed beyond the middle from the costal to the inner margin by a very faint broken black line, which in some specimens is entirely wanting, the fringe white; secondaries smoky-brown, darker on the costal margin and at the apex, the fringe white; the underside of both wings smoky-brown: head and thorax white, the abdomen brownish-white, the palpi and legs blackish-brown, the antennæ black. Expanse 1½ inch.

Hab. MEXICO, Amula in Guerrero (*H. H. Smith*¹).