

3. *Grotella samula*. (Tab. XCIV. fig. 20.)

Grotella samula, Druce, Ann. & Mag. Nat. Hist. (6) xvi. p. 39 (1895) ¹.

Primaries and secondaries citron-yellow, the primaries slightly the darkest, and with the inner margin and a spot at the end of the cell very pale brown; the underside as above, but without the brown markings: head, antennæ, and thorax yellowish (abdomen wanting). Expanse $\frac{3}{4}$ inch.

Hab. MEXICO, Tonalapa in Guerrero (*H. H. Smith* ¹).

One specimen.

ADISURA (I. p. 301).

2. *Adisura grandis*. (Tab. XCIV. fig. 21.)

Adisura grandis, Druce, P. Z. S. 1890, p. 516 ¹.

Primaries bright rose-pink, with a narrow cream-coloured line extending down the middle of the wing from the base to the outer margin, the fringe cream-colour; secondaries pure silky-white, with the fringe white; the underside silky-white, that of the primaries dusky from the base to beyond the middle: head and thorax pink, the base of the latter and the abdomen white, the legs pinkish-white, the antennæ pale pinkish-brown. Expanse $1\frac{1}{2}$ inch.

Hab. MEXICO, Lake Chapala in Jalisco (*Richardson* ¹).

One worn specimen.

ANARTA (to follow the genus *Adisura*, I. p. 301).

Anarta, Hübner. Verz. bek. Schmett. p. 220 (1816); Walker, Cat. xi. p. 696.

1. *Anarta agonax*. (Tab. XCIV. fig. 22.)

Anarta agonax, Druce, P. Z. S. 1890, p. 515 ¹.

Primaries very dark brown, almost black, with a large square white spot at the end of the cell, beyond which the wing is crossed from the costal to the inner margin by a pale yellowish-brown band, this latter being somewhat dentated on the outer side, the fringe pale yellowish-brown, except in the middle, where it is black; secondaries black, broadly white from the base to about the middle of the inner margin, the fringe white; the underside of both wings black, with the white spots as above: head, thorax, antennæ, and legs black. Expanse $\frac{3}{4}$ inch.

Hab. MEXICO, Jalapa (*Höge* ¹).

PANEMERIA (to follow the genus *Anarta*).

Panemeria, Hübner, Verz. bek. Schmett. p. 262 (1816); Walker, Cat. xi. p. 705.

1. *Panemeria rhea*. (Tab. XCIV. fig. 23.)

Panemeria rhea, Druce, Ann. & Mag. Nat. Hist. (6) xiii. p. 360 (1894) ¹.

Primaries dark brown, with a narrow submarginal grey line extending from the apex to the anal angle, the fringe dark brown; secondaries black, crossed about the middle from the costal margin almost to the inner margin by a wide orange-yellow band, the fringe yellowish-brown: head, thorax, and abdomen black. Expanse 1 inch.

Hab. MEXICO, Durango city (*Becker* ¹).