

AGROPHILA (I. p. 301).

8. **Agrophila anæa**, sp. n. (Tab. XCIV. fig. 24.)

Primaries with the basal half fawn-colour, the outer half very dark purplish-brown, marked with some very faint lines near the apex and along the outer margin, the fringe very dark brown; secondaries dark chrome-yellow, the costal margin edged with black, the fringe yellowish-brown: head, tegulæ, and thorax fawn-colour, the abdomen chrome-yellow, the antennæ brown, the legs greyish-brown. Expanse $\frac{1}{2}$ inch.

Hab. MEXICO, Atoyac in Vera Cruz (*H. H. Smith*), Temax in Northern Yucatan (*Gaumer*); GUATEMALA, San Gerónimo (*Champion*).

A specimen from Atoyac is figured.

9. **Agrophila luda**, sp. n. (Tab. XCIV. fig. 25.)

Primaries with the base and part of the costal margin grey, the outer half of the wing blackish-brown; secondaries dark brown; the fringe of both wings dark brown: head, thorax, and abdomen black, the legs greyish-black. Expanse $\frac{1}{2}$ inch.

Hab. MEXICO, Jalapa (*M. Trujillo*).

One worn specimen.

10. **Agrophila orcidia**, sp. n. (Tab. XCIV. fig. 26.)

Primaries brownish-fawn-colour, dark brown at the base and along the outer margin, the fringe also dark brown; secondaries chrome-yellow, shaded with brown, the costal margin narrowly edged with black, the fringe black: head, antennæ, thorax, and abdomen blackish-brown. Expanse $\frac{1}{2}$ inch.

Hab. MEXICO, Omilteme in Guerrero, Teapa in Tabasco (*H. H. Smith*).

This species varies considerably in size. We figure a specimen from Teapa.

11. **Agrophila mata**, sp. n. (Tab. XCIV. fig. 27.)

Primaries greenish-brown, thickly irrorated with yellow scales, with three yellowish-white marks on the costal margin and a fine yellow streak on the inner margin near the anal angle, the marginal line yellowish-white, the fringe dark brown; secondaries brown, paler at the base: head, thorax, and abdomen greenish-brown. Expanse $\frac{7}{10}$ inch.

Hab. MEXICO, Amula in Guerrero, Cuernavaca in Morelos (*H. H. Smith*).

Two specimens. The one from Amula is figured.

METOPONIA (I. p. 303).

4. **Metoponia umbria**, sp. n. (Tab. XCIV. fig. 28.)

Primaries dark brown, paler along the outer margin, and crossed about the middle from the costal to the inner margin by two very indistinct darker brown bands, the fringe brown; secondaries brown, paler at the base, the fringe yellowish-brown: head, antennæ, thorax, and abdomen dark brown. Expanse 1 inch.

Hab. MEXICO, Cuernavaca in Morelos (*H. H. Smith*); PANAMA, Volcan de Chiriqui (*Champion*).

Two specimens. The one from the Volcan de Chiriqui is figured.