

with several dark brown markings near the apex and along the outer margin; the fringe of the primaries dark brown, that of the secondaries yellowish-brown: head, antennæ, thorax, abdomen, and legs dark reddish-brown. Expanse 1 inch.

Hab. PANAMA, Bugaba (*Champion*).

One specimen.

7. **Drobeta debora**, sp. n. (Tab. XCV. fig. 9, ♂.)

Male. Primaries pale pinkish-fawn-colour, crossed about the middle from the costal to the inner margin by a dark brown line, with an indistinct black elongated spot at the end of the cell; secondaries creamy-white, slightly dusky at the apex: head, antennæ, and thorax fawn-colour, the abdomen brownish-white. Expanse 1 inch.

Hab. COSTA RICA, Candelaria Mts. (*Underwood*).

Two males.

8. **Drobeta (?) yeba**, sp. n. (Tab. XCV. fig. 10, ♂.)

Male. Primaries shaded with dark and light brown, and crossed from the costal to the inner margin by two indistinct brown lines, a black dot at the end of the cell; secondaries semihyaline dusky brown, darker at the apex and along the outer margin, the fringe dark brown: head, collar, and tegulæ yellowish-brown, the thorax and abdomen darker brown, the antennæ and legs brown. Expanse $1\frac{1}{10}$ inch.

Hab. MEXICO, Teapa in Tabasco (*H. H. Smith*); PANAMA, Volcan de Chiriqui (*Champion*).

We figure the specimen from the Volcan de Chiriqui.

9. **Drobeta attina**, sp. n. (Tab. XCV. fig. 11, ♀.)

Female. Primaries pale greyish-brown, thickly irrorated with white scales, a wide dark brown curved band crossing the wing from the costal margin to the middle of the outer margin; secondaries brown, edged with small black spots: head and thorax greyish-brown, the antennæ and abdomen pale brown. Expanse $1\frac{4}{10}$ inch.

Hab. PANAMA, Volcan de Chiriqui 3000 feet (*Champion*).

One specimen.

10. **Drobeta leada**, sp. n. (Tab. XCV. fig. 12, ♂.)

Male. Primaries greyish-brown from the base to beyond the middle, the apex greyish-brown, the other portion of the wing reddish-brown, the marginal line pale yellowish-brown, the fringe reddish-brown; secondaries reddish-brown, paler at the base, with a darker brown streak at the end of the cell: head, collar, tegulæ, and thorax greyish-brown, the antennæ and abdomen reddish-brown. Expanse $1\frac{1}{4}$ inch.

Hab. GUATEMALA, Volcan de Atitlan 2500 feet (*Champion*).

One specimen.

PSEUDINA (I. p. 311).

2. **Pseudina chuza**, sp. n. (Tab. XCV. fig. 13, ♀.)

Female. Primaries dark greyish-brown, with a submarginal darker brown band crossing the wing from the apex to the inner margin, two spots near the end of the cell and some markings on the costal margin near the base darker brown, the fringe dark brown; secondaries pale brown, paler at the base: head, collar,