

tegulae, and thorax greyish-brown, the abdomen darker brown, the antennae and legs brown. Expanse  $1\frac{1}{4}$  inch.

*Hab.* MEXICO, Cuernavaca in Morelos (*H. H. Smith*).

One specimen.

### EUSTROTIA (I. p. 311).

#### 13. *Eustrotia decorata*, sp. n. (Tab. XCV. fig. 14, ♂.)

*Male.* Primaries pale primrose-yellow, crossed from the costal to the inner margin by two pale brown bands—the first close to the base, the second beyond the middle,—and with a pale brown spot on the costal margin between them, the apex brown, the fringe greyish; secondaries brownish-white: head, collar, and tegulae pale primrose-yellow, the thorax and abdomen pale brown. Expanse  $\frac{3}{4}$  inch.

*Hab.* MEXICO, Northern Sonora (*Morrison*).

One specimen.

#### 14. *Eustrotia devara*, sp. n. (Tab. XCV. fig. 15.)

Primaries reddish-brown, crossed from the costal to the inner margin by several very indistinct darker waved brown lines, and with a square-shaped black spot at the end of the cell, a minute black dot beyond it, and two yellowish-white streaks on the costal margin—the first just above the end of the cell, the second at the apex,—the fringe dark brown; secondaries pale dusky brown, the fringe pale brown: head, thorax, abdomen, and legs brown. Expanse  $\frac{3}{4}$  inch.

*Hab.* GUATEMALA, Pantaleon 1700 feet (*Champion*); PANAMA, Volcan de Chiriqui 3000 feet (*Champion*).

Two specimens. The one from the Volcan de Chiriqui is figured.

#### 15. *Eustrotia bettina*, sp. n. (Tab. XCV. fig. 16, ♀.)

*Female.* Primaries and secondaries dark brown; the primaries crossed about the middle from the costal to the inner margin by a darker brown band, and with a blackish-brown spot near the apex: head, antennae, thorax, abdomen, and legs dark brown. Expanse  $\frac{3}{4}$  inch.

*Hab.* MEXICO, Jalapa (*M. Trujillo*).

One specimen.

#### 16. *Eustrotia besidia*, sp. n. (Tab. XCV. fig. 17.)

Primaries and secondaries dark brown; the primaries crossed from the costal to the inner margin by two darker brown waved lines—the first near the base, the second beyond the middle,—and with a black spot at the end of the cell and another at the apex; the fringe brown: head, antennae, thorax, abdomen, and legs dark brown. Expanse  $\frac{3}{4}$  inch.

*Hab.* MEXICO, Amula in Guerrero 6000 feet (*H. H. Smith*).

One specimen, probably a female.

#### 17. *Eustrotia cincta*, sp. n. (Tab. XCV. fig. 20, ♂.)

*Male.* Primaries black, with several very minute yellowish spots along the outer margin; secondaries blackish-brown, paler at the base, and with a small black spot at the end of the cell; the fringe black: head, antennae, thorax, and abdomen black, the underside of the latter and the legs pale brown. Expanse  $\frac{3}{4}$  inch.

*Hab.* GUATEMALA, Senahu in Vera Paz (*Champion*).

One specimen.