

# THALPOCHARES (I. p. 314).

## 7. *Thalpochares lagore*. (Tab. XCV. fig. 18.)

*Thalpochares lagore*, Druce, P. Z. S. 1890, p. 517<sup>1</sup>.

Primaries dark brown from the base to the middle, and pale purplish-brown thence to the apex, with a narrow yellow line crossing the wing about the middle from the costal to the inner margin, a black dot at the apex, edged with yellow on the upperside, and three minute yellow streaks on the costal margin near the apex, the marginal line yellow, with very minute black dots on the outer edge, the fringe brown; secondaries uniformly dark brown: head and front of the thorax yellowish-brown, the rest of the thorax, the abdomen, antennæ, and legs dark brown. Expanse  $\frac{3}{4}$  inch.

*Hab.* MEXICO, Teapa in Tabasco (*H. H. Smith*<sup>1</sup>); PANAMA, Chiriqui (*Ribbe, in mus. Staudinger*<sup>1</sup>).

A small dull-coloured species, allied to *T. hippotes*, Druce.

## 8. *Thalpochares laronia*. (Tab. XCV. fig. 19.)

*Thalpochares laronia*, Druce, P. Z. S. 1890, p. 516<sup>1</sup>.

Primaries very pale fawn-colour, crossed by three silvery-white bands, the second band broken into two, forming an oval spot at the end of the cell, the outer margin banded with silvery-white, the space between the first and third bands shaded with bright pink, the fringe white; secondaries white, clouded with very pale fawn-colour: head, thorax, and abdomen pale fawn-colour, the base of the latter whitish, the antennæ and legs pale brown. Expanse 1 inch.

*Hab.* MEXICO, Tierra Colorada in Guerrero 2000 feet (*H. H. Smith*<sup>1</sup>).

One specimen, captured by Mr. Smith in October 1888; it seems to be allied to the European *T. rosea*.

## 9. *Thalpochares darena*, sp. n. (Tab. XCV. fig. 21, ♂.)

*Male.* Primaries very pale brown, crossed about the middle from the costal to the inner margin by a wide dark brown band, which is edged on each side with a whitish line, and with a small dark brown spot near the apex and a very indistinct waved submarginal line extending from the apex to the anal angle, the fringe pale brown; secondaries cream-colour, clouded with pale brown at the apex and round the outer margin, the fringe white: head, antennæ, thorax, collar, and tegulæ pale brown, the abdomen and legs brownish-white. Expanse  $\frac{3}{4}$  inch.

*Hab.* MEXICO, Presidio in Mazatlan (*Forrer*).

Several specimens.

## 10. *Thalpochares* — ?

*Hab.* MEXICO, Jalapa (*M. Trujillo*).

Several specimens, all in very bad condition.

## 11. *Thalpochares laverna*, sp. n. (Tab. XCV. fig. 22, ♂.)

*Male.* Primaries dark brown, crossed from the costal to the inner margin, near the base and beyond the middle, by two very fine, indistinct, lead-coloured lines, and with a large black spot on the costal margin close to the apex, the outer margin broadly bordered with pale brown, the fringe pale brown; secondaries