

inner margin, below which is a reddish-brown spot, a greyish-white line crossing from the apex to the inner margin, and on the outer side of this line is a large subtriangular reddish-brown marking, the point of which just crosses the white line, the fringe pale brown; secondaries pale greyish-brown, darker round the outer margin, the fringe pale brown: head, thorax, abdomen, and legs pale greyish-brown, the antennæ dark brown. Expanse  $1\frac{1}{2}$  inch.

*Hab.* MEXICO, Jalapa (*M. Trujillo*<sup>1</sup>).

One specimen.

15 (B). **Plusia masoni**. (Tab. XCVI. fig. 12.)

*Plusia masoni*, Schaus, Trans. Am. Ent. Soc. xxi. p. 238 (1894)<sup>1</sup>.

*Hab.* MEXICO, Jalapa (*coll. Schaus*<sup>1</sup>, *M. Trujillo*), Orizaba (*Boucard, in mus. D.*).

We figure a specimen of this species from Jalapa.

**Plusia abrota** (I. p. 333).

To the locality given, add:—GUATEMALA, Guatemala city (*Rodriguez*).

THYRIA (I. p. 333).

**Thyria ditissima** (I. p. 333).

To the localities given, add:—BRITISH HONDURAS, Belize (*Sir Alfred Moloney, in mus. D.*).

PLUSIODONTA (to follow the genus *Thyria*, I. p. 333).

*Plusiodonta*, Guenée, Sp. gén. des Lép. vi. p. 359 (1852); Walker, Cat. xii. p. 937.

1. **Plusiodonta thomæ**.

*Plusiodonta thomæ*, Guen. Sp. gén. des Lép. vi. p. 360<sup>1</sup>; Walk. Cat. xii. p. 937<sup>2</sup>.

*Hab.* PANAMA, Chiriqui (*Ribbe*).—ANTILLES, St. Thomas<sup>1 2</sup>.

One specimen of this species has been seen from Chiriqui.

2. **Plusiodonta effulgens**.

*Plusiodonta effulgens*, H. Edwards, Papilio, iv. p. 79 (1884)<sup>1</sup>.

*Hab.* MEXICO, State of Vera Cruz (*Schaus*<sup>1</sup>).

DIAMUNA (I. p. 333).

The genera *Diamuna* and *Clitis* (I. p. 335), included by me in the Noctuidæ, under the subfamily Calpinæ, belong, as already noted, to the family Agaristidæ\*.

\* Cf. *anted*, i. pp. 34–36; ii. pp. 326–329.