

## YPSIA (I. p. 346).

**Ypsia lineata** (I. p. 346).

To the localities given, add:—MEXICO, Jalapa (*M. Trujillo*).

## PHOSPHILA (I. p. 348).

**Phosphila tatosoma** (I. p. 348).

To the localities given, add:—BRITISH HONDURAS, Belize (*Sir Alfred Moloney, in mus. D.*).

## CŒNIPETA (I. p. 349).

**Cœnipeta medina** (I. p. 351).

To the localities given, add:—BRITISH HONDURAS, Belize (*Sir Alfred Moloney, in mus. D.*).

## HYPOGRAMMA (I. p. 351).

**Hypogramma damonia** (I. p. 351).

To the localities given, add:—GUATEMALA, Guatemala city (*Rodriguez, in mus. D.*).

**Hypogramma suttea** (I. p. 352).

To the localities given, add:—COSTA RICA, Candelaria Mts. (*Underwood*).

GADIRTHA (to follow the genus *Decelea*, I. p. 352).

*Gadirtha*, Walker, Cat. xiii. p. 1102 (1857).

1. **Gadirtha candezei**, sp. n. (Tab. XCVI. fig. 17.)

Primaries broadly bluish-black at the base, the costal half of the wing brown flecked with darker brown, the inner half mottled with greyish-black, the fringe alternately black and brown, a brown spot at the anal angle; secondaries greyish-white, broadly shaded with dark brown at the apex and round the outer margin, the fringe greyish-brown: head, antennæ, and palpi black, the collar, tegulæ, and thorax greyish-brown, the abdomen blackish-brown. Expanse  $2\frac{1}{4}$  inches.

*Hab.* GUATEMALA (*Rodriguez, in mus. D.*).

I am indebted to M. L. Candèze for a specimen of this fine species.

2. **Gadirtha (?) leena**, sp. n. (Tab. XCVI. fig. 19, ♂.)

*Male.* Primaries pale brown, mottled with darker brown about the middle and along the outer margin, and with a dark brown mark on the costal margin close to the base and a large similarly-coloured Y-shaped mark near the apex, the fringe pale brown; secondaries pale brown, with a row of very minute yellowish spots round the margin: head, antennæ, thorax, and abdomen brown. Expanse  $1\frac{3}{4}$  inch.

*Hab.* PANAMA, Bugaba (*Champion*).

One specimen.