

PROMETOPUS (to follow the genus *Gadirtha*).

*Prometopus*, Guenée, Sp. gén. des Lép. v. p. 38 (1852); Walker, Cat. xiii. p. 1119.

1. *Prometopus ordinarius*, sp. n. (Tab. XCVI. fig. 20.)

Primaries blackish-brown, almost black at the base and along part of the inner margin, with a greyish spot at the end of the cell, another one below it, and a few black spots along the outer margin; secondaries brownish-white, dark brown from the apex to the anal angle, the fringe pale brown: head, antennæ, collar, and tegulæ dark brown, the thorax and abdomen pale brown, the anal tuft yellowish. Expanse  $1\frac{1}{2}$  inch.

*Hab.* MEXICO, Jalapa (*M. Trujillo*); COSTA RICA, Candelaria Mts. (*Underwood*); PANAMA, Volcan de Chiriqui 2000 to 3000 feet (*Champion*).

There are specimens of this species in the British Museum labelled with the above name, but I am unable to find any published description of the insect. An example from the Volcan de Chiriqui is figured.

TRISULOIDES (to follow the genus *Prometopus*).

*Trisuloides*, Butler, Ann. & Mag. Nat. Hist. (5) vii. p. 36 (1881).

This genus was referred to the Liparidæ by Mr. Butler, who notes the resemblance of his Indian type to the genera *Chrysorithrum*, *Allotria*, and *Catocala*.

1. *Trisuloides pata*. (Tab. XCVI. fig. 21.)

*Trisulodes pata*, Druce, Ann. & Mag. Nat. Hist. (6) xiii. p. 362 (1894)<sup>1</sup>.

Primaries greyish-brown, crossed from the costal to the inner margin by several waved black lines, with a round grey spot in the cell, a pale grey band beyond it, this being widened towards the costal margin, and a wide black streak near the anal angle, the fringe alternately black and grey; secondaries pale yellow, broadly bordered with blackish-brown from the apex to the anal angle: head and thorax dark grey, the abdomen black. Expanse 2 inches.

*Hab.* GUATEMALA, in the city (*Rodriguez*<sup>1</sup>).

COCYTODES (to precede the genus *Stictoptera*, I. p. 353).

*Cocytodes*, Guenée, Sp. gén. des Lép. vii. p. 41 (1852); Walker, Cat. xiii. p. 1122.

1. *Cocytodes schneideriana*.

*Phalæna Noctua schneideriana*, Cram. Pap. Exot. iv. p. 37, t. 308. fig. A<sup>1</sup>; Stoll, Suppl. Cram. Pap. Exot. v. p. 109, t. 22. fig. 8 (larva)<sup>2</sup>.

*Letis schneideriana*, Guen. Sp. gén. des Lép. vii. p. 149<sup>3</sup>; Walk. Cat. xiv. p. 1269<sup>4</sup>.

*Achæa schneideriana*, Möschl. Verh. zool.-bot. Ges. Wien, 1880, p. 429<sup>5</sup>.

*Hab.* MEXICO, Orizaba (*Boucard, in mus. D.*), Jalapa (*M. Trujillo*); GUATEMALA, Guatemala city (*Rodriguez*); COSTA RICA, Candelaria Mts. (*Underwood*). — GUIANA, Surinam<sup>1 2 3 4 5</sup>.

Of this species I have seen five specimens from our region. It is very closely allied to the Indian *C. cœrula* and *C. granulata*, Guen., but has the primaries narrower and more pointed at the apex.