

STICTOPTERA (I. p. 353).

Stictoptera fenestra (I. p. 353).

To the localities given, add:—MEXICO, Jalapa (*M. Trujillo*); COSTA RICA, Candelaria Mts. (*Underwood*).

MELIPOTIS (I. p. 354).

Melipotis fasciolaris (I. p. 354).

To the localities given, add:—GUATEMALA, Guatemala city (*Rodriguez*).

Melipotis nigrescens (I. p. 356).

To the localities given, add:—COSTA RICA, Candelaria Mts. (*Underwood*).

Melipotis agrotoides (I. p. 358).

To the localities given, add:—BRITISH HONDURAS, Belize (*Sir Alfred Moloney, in mus. D.*); COSTA RICA, Candelaria Mts. (*Underwood*).

23. **Melipotis inconspicua.** (Tab. XCVI. fig. 22.)

Bolina inconspicua, Schaus, Trans. Am. Ent. Soc. xxi. p. 240 (1894)¹.

Hab. MEXICO, Guadalajara in Jalisco (*Goldsmith, in mus. D.*), Jalapa (*coll. Schaus*¹, *M. Trujillo*); GUATEMALA, Guatemala city (*Rodriguez*).

Varies considerably in colour. The single specimen received from Guatemala is paler than the others, and it also has the apical black dots more distinct. We figure an example from Jalapa.

24. **Melipotis usipetes**, sp. n. (Tab. XCVI. fig. 23.)

Primaries blackish-brown, thickly irrorated with bluish-grey scales, with the costal margin from the base to beyond the middle pale brown and flecked with white scales, a round white dot in the cell, a large oval white spot at the end of the cell extending across the wing almost to the inner margin, and several yellowish-brown spots close to the apex, the fringe blackish-brown; secondaries black, with a central white line extending from the base to about the middle, the fringe black, a white patch about the middle of the outer margin excepted: head, antennæ, collar, tegulæ, and thorax brownish-black, the abdomen greyish-black, the underside of the thorax and abdomen greyish-brown. Expanse $1\frac{3}{4}$ inch.

Hab. MEXICO, Jalapa (*M. Trujillo*); COSTA RICA, Volcan de Irazu 6000 to 7000 feet (*Rogers*), Candelaria Mts. (*Underwood*).

A specimen from Jalapa is figured.

CIRRHOBOLOINA (I. p. 358).

Cirrhopolina deducta (I. p. 358). (Tab. XCVI. fig. 24.)

We now figure our Panama specimen of this species.

Cirrhopolina mexicana (I. p. 359).

To the Mexican localities given, add:—Guadalajara in Jalisco (*Goldsmith, in mus. D.*).