

1. *Walshia calcarata*, sp. n. (Tab. I. fig. 15.)

Antennae brownish, the last eleven or twelve joints pure white. *Palpi* with the median joint enlarged by projecting scales above, and more especially beneath, giving it a triangular form; terminal joint stout, somewhat clothed beneath nearly to the apex; white, dusted with brown, except at the extremity of each joint. *Head* white. *Thorax* greyish fuscous above, whitish behind; tegulae whitish. *Forewings* brown and whitish, mixed in about equal proportions, with a lilac gloss about the paler parts, especially below the fold; a stout, square-ended, tuft of brownish ochreous raised scales below the fold at one-third of its length, a ridge of raised scales of the same colour, but delicately striated with brown, crosses the middle of the wing from costa to dorsum; another raised patch, somewhat tinged with fuscous, occurring at the end of the cell, is succeeded by a pale transverse band with lilac reflections, the central fascia being also preceded and followed by the same whitish ground-colour; three fuscous dots occur at the base of the brownish ochreous cilia below the apex, the tornal cilia tending to greyish. *Exp. al.* 13 mm. *Hindwings* cinereous, dusted with pale brownish; cilia cinereous, tinged with ochreous towards the apex. *Abdomen* (missing). *Legs* brownish, the hind tibiae obliquely banded with white at the base of the first pair of spurs; second pair of spurs densely clothed with a triangular tuft of white scales above; tarsi slightly annulate with white.

Type ♂ (65539) Mus. Wlsm. (*Godm-Salv. Coll.*).

Hab. MEXICO: TABASCO: Teapa, III. (*H. H. Smith*). Unique.

The clothing of the spurs of the hind tibiae distinguishes *calcarella* from *amorphella* Clms., and *detracta* Wlsm.

2. *Walshia detracta*, sp. n. (Tab. I. fig. 14.)

Antennae cinereous, annulate with brownish fuscous; the elongate basal joint fuscous above, pale beneath. *Palpi* recurved to the vertex, somewhat coarsely clothed nearly to the apex, terminal joint longer than median; pale cinereous, shaded externally with fuscous nearly to the end of the median joint and above the middle of the terminal joint. *Head* greyish fuscous above, face steel-whitish. *Thorax* dark greyish fuscous. *Forewings* dark greyish fuscous to the end of the basal third, with a large concurrent pattern of raised scales, tinged with chestnut-brown beyond the base, especially about the middle of the costa and at the apex, with numerous tufts of raised scales distributed nearly to the apex; cilia brownish cinereous, greyish fuscous at the extreme apex. *Exp. al.* 15 mm. *Hindwings* greyish brown; cilia 6, brownish cinereous. *Legs* greyish fuscous, with whitish bands at the base of each pair of spurs.

Type ♀ (65538) Mus. Wlsm. (*Godm-Salv. Coll.*).

Hab. MEXICO: GUERRERO: Amula, 6000 ft., VIII. (*H. H. Smith*). Unique.

LAVERNA Crt.

=† *LARUNA* Chmb., *laps. cal.*; =* *MOMPHA* (Hb.) HS., *nec* Stph.; = *WILSONIA* Clms.

Type 1: *Laverna ochraceella* Crt. (Crt. 1839).

LAVERNA Crt. Br. Ent. 16 expl. Pl. 735 (1839); Stn. List Br. An. BM. XVI. Lp. 3 137-9 (1854); Ins. Br. Lp-Tin. 225, 235-40 Pf. 7·8^{a-c} (1854); Frey Tin-Pter. Schweiz 276-83 (1856); Lin. Ent. 14 180-205 (1860); Wkr. Cat. Lp. BM. 30 882-5 (1864); Stn. NH. Tin. 11 126-49 Pl. 4-6 (1870); Stgr-Wk. Cat. Lp. Eur. 319 sp. 2569-88 (1871); Clms-Stn. Tin. N-Am. 130-1 (1872); [† *LARUNA* Chmb. Can. Ent. 5 229 (1873): 6 49, 73 (1874)]; Hnmn. Schm. Deutsch. Tin. 416-20, Tbl. 8, 53-4 (1876); Z. Hor. Soc. Ent. Ross. 13 420-7 (1877); Chmb. Bull. US. GG. Surv. 4 151-2 (1878); Snln. Vlind. Ned. Micr. 422, 852-60 Pf. 12·419-22 (1882); Wlsm. Tr. Ent. Soc. Lond. 1891 117-8 Pf. 5-6·54-5 (1891); Riley, Smith's List Lp.