

ARISTOTELIA Hb.

= CHRYSOPORA Clms. (= § *NOMIA* Clms.) ; = NANNODIA Hnmn. ; = § *ERGATIS* Hnmn. ; = XYSTOPHORA Hnmn-Wk. (= § *DORYPHORA* Hnmn.) ; = MONOCHROA Hnmn. ; = § *LAMPROTES* Hnmn. ; = EUCATOPTUS Wlsm.

Type 1 : *Tinea decurtella* Hb. Pf. 45 · 311 (Hb. 1826).

ARISTOTELIA Hb. Verz. Schm. 424 sp. 4133 (1826) ; Wlsm. Ent. Mo. Mag. 31 42 (1895) ; Meyr. HB. Br. Lp. 569, 572-7 no. 4 sp. 1-21 tf. 572 (1895) ; Wlsm. Pr. Z. Soc. Lond. 1897 63-8 no. 10 sp. 21-32 (1897) ; Busck Bull. US. Nat. Mus. 52 498-9 sp. 5575-91 (1902) ; Pr. US. Nat. Mus. 25 773, 794-804 Pf. 29 · 11 (1903) : 27 755-7 (1904) : 30 724-5 (1906) ; Pr. Ent. Soc. Wash. 11 87 (1909) ; Meyr. Pr. Lin. Soc. NSW. 29 257, 286-92 no. 7 sp. 51-60 (1904) ; Jr. Bomb. NH. Soc. 17 138 (1906) : 18 437-9 (1908) ; Pr. Z. Soc. Lond. 1908 724 (1908) ; Ann. S-Afr. Mus. 5 369-70 (1909) ; Wlsm. Fn. Haw. 1 478-82, 744 no. 5 sp. 11-19 Pf. 13 · 17-25 (1907) ; Pr. Z. Soc. Lond. 1907 930-1 no. 14 sp. 25-6 (1908).

Type 2 : *Nomia lingulacella* Clms. (Clms. 1860).

CHRYSOPORA Clms. = § *NOMIA* Clms. Pr. Ac. Nat. Sc. Phil. 12 167-8 (1860) : Clms-Stn. Tin. N-Am. 123-4 (1872). CHRYSOPORA Clms. (nn.) Pr. Ac. Nat. Sc. Phil. 12 362 (1860) : Clms-Stn. Tin. N-Am. 123, 158 (1872) ; Wlsm. Ent. Mo. Mag. 31 42 (1895) ; Stgr-Rbl. Cat. Lp. Pal. 2 156 no. 319 sp. 2894-7 (1901) ; Busck Bull. US. Nat. Mus. 52 497-8 sp. 5572-3 (1902) ; Pr. US. Nat. Mus. 25 773, 792-3 Pf. 29 · 9 (1903) ; Wlsm. Pr. Z. Soc. Lond. 1907 931-2 no. 15 sp. 27 Pf. 51 · 7 (1908).

In *Chrysopora lingulacella* Clms., as also in *Nannodia stipella* Hb. and *hermannella* F., vein HW : 6, though weak, is remote from 7 and parallel with the fold ; the discoidal is traceable, and Meyrick was justified in referring the types of *Nannodia* Hnmn. to *Aristotelia* Hb.

Type 3 : *Tinea stipella* Hb.

NANNODIA Hnmn. Schm. Deutsch. Tin. 190, 284-5 no. 426 sp. 423-4 (1870), Tbl. 5, 41 no. 426 (1876) ; Stn. Ent. Mo. Mag. 7 168 (1871) ; Snln. Vlind. Nederl. Micr. 432, 690-2 no. 53 sp. 1-2 Pf. 9 · 286 (1882).

Type 4 : *Oecophora brizella* Tr.

§ *ERGATIS* Hnmn. Schm. Deutsch. Tin. 190, 295-8 no. 432 sp. 436-41 (1870), Tbl. 5, 42 no. 432 (1876) ; Stn. Ent. Mo. Mag. 7 169-70 (1871) ; Snln. Vlind. Nederl. Micr. 431, 681-4 no. 50 sp. 1-3 Pf. 9 · 289 (1882).

Type 5 : *Anacamptis pulveratella* HS.

XYSTOPHORA Hnmn-Wk. = § *DORYPHORA* Hnmn. Schm. Deutsch. Tin. 190, 298-308 no. 433 sp. 442-60 (1870) ; Stn. Ent. Mo. Mag. 7 170 (1871) ; Z. Hor. Soc. Ent. Ross. 13 359-61 no. 19 (1877). XYSTOPHORA Hnmn-Wk. (nn.) Tbl. 6 no. 433, 42 no. 433 (1876) ; Snln. Vlind. Nederl. Micr. 431-2, 684-8 no. 51 sp. 1-5 Pf. 9 · 287, 12 · 404^{a-b} (1882) ; Drnt. Bull. Soc. Ent. Fr. 1889 p. ccxxiv (1889) ; Wlsm. Ent. Mo. Mag. 31 42 (1895) ; Stgr-Rbl. Cat. Lp. Pal. 2 152-3 no. 310 sp. 2786-2824 (1901).