

Type 6: *Tinea tenebrella* Hb. (Hnmn. 1870).

MONOCHROA Hnmn. Schm. Deutsch. Tin. 190, 308-9 no. 434 sp. 461 (1870), Tbl. 6, 43 no. 434 (1876); Stn. Ent. Mo. Mag. 7 170 (1871).

Type 7: *Tinea atrella* (Hb.?) Hw.

§ LAMPROTES Hnmn. Schm. Deutsch. Tin. 190, 309-11 no. 435 sp. 462-6 (1870), Tbl. 6, 43 no. 435 (1876); Stn. Mo. Mag. 7 170-1 (1871).

Type 8: *Eucatoptus penicillata* Wlsm. (Wlsm. 1897).

EUCATOPTUS Wlsm. Pr. Z. Soc. Lond. 1897 69-71 no. 11 sp. 33-6 (1897); Busck Pr. US. Nat. Mus. 25 794-5 (1903).

## 1. *Aristotelia corallina*, sp. n.

*Antennae* blackish. *Palpi* with the median joint somewhat expanded outwardly; cream-whitish, banded externally with a brownish shade; terminal joint whitish, with two blackish bands. *Head* rosy cream-colour; *Thorax* the same, but with a stronger reddish tinge, the tegulae blackish. *Forewings* blackish, the dorsum rich rosy reddish, this colour diffused upward along the termen and over the terminal cilia through which runs a somewhat obscurely reduplicated dark shade-line; terminal cilia blackish, preceded by three small rosy reddish costal spots. *Exp. al.* 12 mm. *Hindwings* dark grey; cilia brownish cinereous. *Abdomen* brownish cinereous. *Legs* brownish cinereous, broadly banded with fuscous externally on each joint.

*Type* ♂ (65760) Mus. Wlsm. (*Godm-Salv. Coll.*). [*PT.* (65762) US. Nat. Mus.]

*Hab.* MEXICO: GUERRERO: Amula, 6000 ft., VIII-IX. (*H. H. Smith*). Three specimens.

The rosy cream of the head, thorax, and dorsum, contrasted with the blackish forewings, renders this species somewhat conspicuous.

## 2. *Aristotelia howardi*, sp. n. (Tab. I. fig. 21.)

*Antennae* blackish, with a series of elongate white spots along the upper side; basal joint reddish ochreous. *Palpi* (broken, the basal joint indicating that they were pale ochreous, or whitish, on their inner sides, and banded, or shaded, with brown, or blackish, externally). *Head* rosy ochreous. *Thorax* mottled with reddish brown and ochreous, with a grey median patch. *Forewings* milk-white, tinged with rosy pink along the costa, except at the base and along the dorsum beyond the middle; at the base is a tawny-red patch, wider on the dorsum than on the costa, narrowly darkened to black along its oblique outer margin; after an oblique fascia of the white ground-colour, which is produced outward along the dorsum, where it is tinged with bright ochreous, a rather oblique, mixed black and reddish grey streak descends from the costa at one-fourth, crossing the cell, and bent outward and upward at its lower extremity enclosing above it a patch of the ground-colour profusely stippled with pale rosy grey, bounded on its outer side by a darker, tawny reddish grey quadrangular costal spot, extending downward to an almost circular ochreous spot at the end of the cell; the terminal area is thickly sprinkled with rust-brown, and contains some black spots, notably two on the costa before the apex, in strong contrast to the rosy pink by which they are separated and partially surrounded; cilia whitish cinereous, with three pale greyish wavy shade-lines running through them; underside fuscous, with the middle of the costa narrowly touched with white, a rosy white streak running through the costal cilia. *Exp. al.* 13.5 mm. *Hindwings* shining, steel-grey, with a bright rosy iridescence; cilia pale brownish cinereous. *Abdomen*