

6. *Aristotelia pyrodercia*, sp. n.

Antennae delicately annulated with black and white. *Palpi* yellowish white, the median joint twice annulated with brassy brown; terminal joint longer than the median, with three blackish annulations, widening in succession from the base. *Head* pale buff-brown. *Thorax* pale buff-brown, with a rather shining steel-brown median streak. *Forewings* yellowish brown, crossed by oblique whitish streaks mixed with shining steel-like metallic scales, some minute blackish scales intermixed along their margins; a slender steely line, commencing at the base of the costa, crosses the fold obliquely outward to the dorsum, beyond it is a broader whitish costal streak, also crossing the fold obliquely outward, but scarcely reaching the dorsum; a third steely streak, from the costa before the middle, tending obliquely outward, is enlarged at the lower end of the cell and reverts to the middle of the dorsum, it is preceded by a pale plical spot margined on its inner side by black scales; beyond this, about the middle of the costa, an oblique whitish streak tends outward to the end of the cell where it meets a more conspicuous white streak inverted from the costal cilia, both mixed with pale steel and blackish scales; beyond the last white costal streak the costal cilia are greyish fuscous, with two inverted whitish streaklets at their base, a third shining white streak following the termen at the base of the whitish cilia through which run two dark shade-lines terminating before the tornus where a yellowish white line in the tornal cilia interrupts them; near the base of this is a luminous steely spot, with some minute black scales along the margin. *Exp. al.* 12 mm. *Hindwings* and cilia tawny greyish. *Abdomen* brownish cinereous. *Legs* whitish, banded externally on the tibiae and tarsi with brownish fuscous.

Type ♀ (65774) Mus. Wlsm. (*Godm-Salv. Coll.*). [*PT.* (65775) US. Nat. Mus.]

Hab. MEXICO: GUERRERO: Tierra Colorada, 2000 ft., X. (*H. H. Smith*); Amula, 6000 ft., VIII. (*H. H. Smith*). Three specimens.

This species differs from *hieroglyphica* Wlsm. in the more olivaceous brownish-ochreous ground-colour which lacks the blackish shades, in the basal half especially, and the silvery markings connected with the outer streaks are more conspicuous.

7. *Aristotelia sarcodes*, sp. n. (Tab. I. fig. 23.)

Antennae black, with elongate white spots along the upper side; basal joint white, with a black line beneath and a spot above. *Palpi* white, median joint with two black bands not reaching the upper surface; terminal joint with four black bands of which the outer two only complete its circumference. *Head* pale brownish, mixed with whitish scales. *Thorax* pale rust-brown, with some silvery median scales. *Forewings* yellowish brown, or light rust-brown, with white and shining silvery costal and dorsal streaks, more or less confluent on the wing-middle, and with two, noticeable, rosy flesh-coloured, short length-streaks, one at the upper edge of the outer end of the cell, the other wider, beyond the end of the cell; there is a rather oblique white costal streak, at one-fifth, with a few black scales along its inner edge, its outer half touched with shining silvery; this meets the apex of a short silvery dorsal streak, at about one-fourth, and is followed by a second, shorter, silvery and white costal streak to which there is no corresponding dorsal one; a third, silvery and white costal streak, is scarcely beyond the middle, and is interrupted by the first rosy flesh-coloured streak before joining a longer silvery oblique streak from the middle of the dorsum; beyond this again are four short white costal streaks before the apex, the first two sometimes confluent, and both tipped with silvery, the second rosy flesh-coloured length-streak separating them from a long silvery dorsal streak arising obliquely outward from near the tornus; there are also a few silvery scales on the termen below the outer white costal streak which occupies the apex and apical cilia; terminal cilia whitish, touched with brown on the middle of the termen and at the tornus. *Exp. al.* 10 mm. *Hindwings* dark slaty grey, with a strong rosy iridescence; cilia brownish grey; ♂ with dull ochreous subcostal hair-pencil. *Abdomen* grey. *Legs* white, hind tibiae and tarsi with black bands across their outer sides.

Type ♂ (5842, Wlsm. Det. 1908) US. Nat. Mus.

Hab. PANAMA: CANAL ZONE: Tabernilla, 1907 (*A. Busck*). Unique.