

4. *Anacamopsis conistica*, sp. n.

Antennae brownish cinereous, dusted with black above. *Palpi* dull white, the median joint distinctly dark brown from its base on the outer side to beyond half its length, the dark brown abruptly cut off; the terminal joint brown on its outer side and around its extreme apex. *Head* greyish brown, face paler. *Thorax* slightly darker than the head. *Forewings* brownish cinereous, much suffused with fuscous, becoming tawny towards the apex and termen, a small black spot at the extreme base of the costa; four very indistinct fuscous spots, two on the fold, and two on the cell, of which the first discal is beyond the second plical; a dull brownish ochreous spot, at the commencement of the costal cilia, connected with an even more obscure opposite dorsal spot by a scarcely distinguishable line of pale scales, outwardly bowed among the fuscous suffusion; a few blackish scale-spots around the apex and termen, at the base of the greyish fuscous cilia which are minutely sprinkled along their middle with whitish cinereous, and faintly tipped with brownish ochreous; the apex is not greatly depressed, but somewhat obtuse. *Exp. al.* 14–15 mm. *Hindwings* scarcely broader than the forewings, trapezoidal, not indented; greyish brown; cilia tawny brownish grey. *Abdomen* brownish fuscous. *Legs* cinereous, suffused on their outer sides with tawny fuscous.

Type ♂ (35711, Sonora) Mus. Wlsm. [PTT. (35712–3) US. Nat. Mus.]

Hab. MEXICO: SONORA: 1883 (*H. K. Morrison*): DURANGO: Presidio (*Forrer*). Three specimens.

This species is very close to 5702 *rhoifructella* Clms., from some varieties of which it may possibly be undistinguishable by the colour or markings of the wings, but so far as I am at present aware the palpi of *rhoifructella* are smoother, less densely clothed, and suffused at least to the apex of the median joint with the darker shading, which is never abruptly cut off leaving the apex cleanly whitish, as in the Mexican species here described.

5. *Anacamopsis quinquepunctella* Wlsm. (Tab. I. fig. 35.)

Anacamopsis quinquepunctella Wlsm. Pr. Z. Soc. Lond. 1897 80–1 no. 61 (1897)¹.

Type ♂ (65179) Mus. Wlsm. (*Godm-Salv. Coll.*). [PT. (65181) US. Nat. Mus.]

Hab. AMERICA, C-I. Central America—MEXICO: TABASCO: Teapa, III. (*H. H. Smith*). West Indies—GRENADA: Balthazar, windward side, 5–15 IV. (*H. H. Smith*)¹.

A single specimen of this species (65710) was received from Teapa.

6. *Anacamopsis ursula*, sp. n.

Antennae brownish grey. *Palpi* cinereous, the short median joint somewhat widened, shaded with greyish fuscous externally nearly to its apex, terminal joint very long, slender, recurved. *Head* mouse-grey; face shining cinereous. *Thorax* mouse-grey, with greenish and violet iridescence. *Forewings* with the costa straight, the apex depressed and somewhat angular; mouse-grey, sprinkled with greyish fuscous scales, a minute fuscous spot at the extreme base of the costa; from the commencement of the costal cilia an indistinct narrow pale cinereous and somewhat sinuate band, curving outward to below the middle, reverts to the dorsum before the tornus, beyond it the apical and terminal space is cupreous, with a slight iridescent lustre; cilia dull brownish cinereous, a few fuscous scales at their base around the apex. *Exp. al.* 18 mm. *Hindwings* dark brown, somewhat thinly scaled toward base; cilia brownish cinereous. *Abdomen* flattened; greyish brown. *Legs* cinereous, much shaded with greyish brown.

Type ♂ (65746) Mus. Wlsm. (*Godm-Salv. Coll.*). [Type ♀ (65772) US. Nat. Mus.]

Hab. MEXICO: MORELOS: Cuernavaca, VI. (*H. H. Smith*). Two specimens.